antibodies.com

Datasheet for ABIN5013544 anti-Fibrillarin antibody (AA 85-274)

3 Images



Overview

Quantity:	100 μL
Target:	Fibrillarin (FBL)
Binding Specificity:	AA 85-274
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fibrillarin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	FBL (Gly85-Cys274)
lsotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FBL. It has been selected for its ability to recognize FBL in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	


Target:	Fibrillarin (FBL)
Abstract:	FBL Products
Background:	Alternative Names: FIB, FLRN, RNU3IP1, rRNA 2'-O-Methyltransferase Fibrillarin, 34 kDa

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5013544 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details

	nucleolar scleroderma antigen, Histone-glutamine methyltransferase
Pathways:	Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	• Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.


Images



Western Blotting

Image 1. Western Blot; Sample: Human A375 cell lysate; Primary Ab: 3µg/ml Rabbit Anti-Rat FBL Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)

Images



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.

Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue

