

Datasheet for ABIN5013545 anti-Fibrillarin antibody (AA 37-275)

2 Images




Overview

Overview	
Quantity:	100 μL
Target:	Fibrillarin (FBL)
Binding Specificity:	AA 37-275
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Fibrillarin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	FBL (Gly37-Ala275)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FBL. It has been selected for its ability to recognize FBL in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography
Target Details	
Target:	Fibrillarin (FBL)
Abstract:	FBL Products
Background:	Alternative Names: FIB, FLRN, RNU3IP1, rRNA 2'-O-Methyltransferase Fibrillarin, 34 kDa


	nucleolar scleroderma antigen, Histone-glutamine methyltransferase
Pathways:	Ribonucleoside Biosynthetic Process
Application Details	
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded kidney tissue