antibodies -online.com









Go to	D		
	Pron	HCL	nane

Overview	
Quantity:	100 μL
Target:	FLRT2
Binding Specificity:	AA 300-517
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FLRT2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	FLRT2 (Ser300-Ser517)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against FLRT2. It has been selected for its ability to recognize FLRT2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Tarnet Details

Background:	Alternative Names: Leucine-rich repeat transmembrane protein FLRT2, Fibronectin-like domain-
Abstract:	FLRT2 Products
Target:	FLRT2
- Target Details	

containing leucine-rich transmembrane protein 2

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded heart tissue