



[Go to Product page](#)

Datasheet for ABIN5013566
anti-FVT1 antibody (AA 64-271)

2 Images

Overview

| | |
|----------------------|--|
| Quantity: | 100 µL |
| Target: | FVT1 (KDSR) |
| Binding Specificity: | AA 64-271 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This FVT1 antibody is un-conjugated |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

Product Details

| | |
|---------------|--|
| Immunogen: | FVT1 (Arg64-Gly271) |
| Isotype: | IgG |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against FVT1. It has been selected for its ability to recognize FVT1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography |

Target Details

| | |
|-------------------|--|
| Target: | FVT1 (KDSR) |
| Alternative Name: | Follicular Lymphoma Variant Translocation 1 (FVT1) (KDSR Products) |
| Background: | Alternative Names: KDSR, DHSR, SDR35C1, 3-Ketodihydrosphingosine Reductase, 3- |

Target Details

Dehydrosphinganine reductase, Short Chain Dehydrogenase/Reductase Family 35C,Member 1

Application Details

Application Notes:

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only

Handling

Format:

Liquid


Concentration:

Lot specific

Buffer:


PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded testis tissue