

Datasheet for ABIN5013592

anti-GABBR1 antibody (AA 199-570)

3 Images



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	GABBR1
Binding Specificity:	AA 199-570
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GABBR1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunoprecipitation (IP)

Product Details

Purpose:	Polyclonal Antibody to Gamma-Aminobutyric Acid B Receptor 1 (gABBR1)
Immunogen:	RPC934Hu01Recombinant GammaAminobutyric Acid B Receptor 1 (gABBR1)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against gABBR1. It has been selected for its ability to recognize gABBR1 in immunohistochemical staining and western blotting.
Cross-Reactivity:	Mouse, Pig, Rat
Purification:	Antigen-specific affinity chromatography followed by Protein A affinity chromatography

Target Details


Target:	GABBR1
Alternative Name:	gABBR1 (GABBR1 Products)
Background:	GABAB(1e), GABABR1, GABA-BR1, GABBR1-3, GPRC3A, HGB1a, Gamma-aminobutyric acid type B receptor subunit 1
Pathways:	Positive Regulation of Peptide Hormone Secretion , cAMP Metabolic Process

Application Details

Application Notes:	Western blotting: 0.01-2 µg/mL, Immunohistochemistry: 5-20 µg/mL, Immunocytochemistry: 5-20 µg/mL, Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Preservative:	Sodium azide
Storage:	4 °C, -20 °C
Storage Comment:	Store at 4°C for frequent use. Stored at -20°C in a manual defrost freezer for two year without detectable loss of activity. Avoid repeated freeze-thaw cycles.




Immunohistochemistry

Image 1. Used in DAB staining on formalin fixed paraffin-embedded heart tissue



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Human Liver Tissue