

## Datasheet for ABIN5013695

# anti-HMGB2 antibody (AA 2-187)





Go to Product page

()	ve	rvi	6	W
$\sim$	v C	1 V I	$\sim$	v v

Quantity:	100 μL	
Target:	HMGB2	
Binding Specificity:	AA 2-187	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This HMGB2 antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	HMGB2 (Gly2-Pro187)	
Isotype:	IgG	
Isotype: Specificity:	IgG  The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its ability to recognize HMGB2 in immunohistochemical staining and western blotting.	
	The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its	
Specificity:	The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its ability to recognize HMGB2 in immunohistochemical staining and western blotting.	
Specificity:  Purification:	The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its ability to recognize HMGB2 in immunohistochemical staining and western blotting.	
Specificity:  Purification:  Target Details	The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its ability to recognize HMGB2 in immunohistochemical staining and western blotting.  Antigen-specific affinity chromatography	
Specificity:  Purification:  Target Details  Target:	The antibody is a rabbit polyclonal antibody raised against HMGB2. It has been selected for its ability to recognize HMGB2 in immunohistochemical staining and western blotting.  Antigen-specific affinity chromatography  HMGB2	

#### **Target Details**

Pathways:
-----------

Cellular Response to Molecule of Bacterial Origin

### **Application Details**

#### Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

#### Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only

#### Handling

Format:	Liquid
Concentration:	Lot specific

Buffer:


PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

#### **Images**




#### **Western Blotting**

Image 1. Western Blot; Sample: Human 293T cell lysate; Primary Ab: 1μg/ml Rabbit Anti-Human HMGB2 Antibody Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit lgG Polyclonal Antibody (Catalog: SAA544Rb19)



#### **Western Blotting**

**Image 2.** Figure. Western Blot; Sample: Recombinant protein.



#### **Immunohistochemistry**

**Image 3.** Used in DAB staining on fromalin fixed paraffinembedded liver tissue