

Datasheet for ABIN5013786  
**anti-IL17RA antibody (AA 617-847)**[Go to Product page](#)

## 3 Images

## Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | IL17RA   |
| Binding Specificity: | AA 617-847   |
| Reactivity:          | Rat  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This IL17RA antibody is un-conjugated  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

## Product Details

|               |  |
|---------------|--|
| Immunogen:    | IL17RA (Gly617-Gly847)   |
| Isotype:      | IgG  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against IL17RA. It has been selected for its ability to recognize IL17RA in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography   |

## Target Details

|             |   |
|-------------|---|
| Target:     | IL17RA  |
| Abstract:   | <a href="#">IL17RA Products</a>   |
| Background: | Alternative Names: CDw217, CD217, IL17R-A, IL17-RA, IL17R-A, IL17R, HIL-17R |

## Target Details

Pathways: [SARS-CoV-2 Protein Interactome](#)

## Application Details

Application Notes:

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment: The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions: For Research Use only


## Handling

Format: Liquid

Concentration: Lot specific


Buffer: PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

## Images




### Immunohistochemistry

**Image 1.** DAB staining on IHC-P; Samples: Rat Brain Tissue



Immunohistochemistry

**Image 2.** Used in DAB staining on formalin fixed paraffin-embedded Kidney tissue



Western Blotting

**Image 3.** Figure. Western Blot; Sample: Recombinant protein.