# antibodies .- online.com







# anti-IVD antibody (AA 1-235)





$\sim$					
( )	VE	۲۱	/1	$\triangle$	Λ

Background:

Quantity:	100 μL	
Target:	IVD	
Binding Specificity:	AA 1-235	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IVD antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
Immunogen:	IVD (Met1-Phe235)	
Isotype:	IgG	
Specificity:	The antibody is a rabbit polyclonal antibody raised against IVD. It has been selected for its ability to recognize IVD in immunohistochemical staining and western blotting.	
Purification:	Antigen-specific affinity chromatography	
Target Details		
Target:	IVD	
Alternative Name:	Isovaleryl Coenzyme A Dehydrogenase (IVD) (IVD Products)	

Alternative Names: ACAD2

Pathways:

Monocarboxylic Acid Catabolic Process

# **Application Details**

# Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

#### Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only


# Handling

Format:	Liquid
Concentration:	Lot specific

Buffer:


PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

### **Images**



# **Western Blotting**

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



# **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded kidney tissue