

# Datasheet for ABIN5013997 anti-NOV antibody (AA 139-357)

## 2 Images




Go to Product page

_				
( )	ve.	rv/	101	Λ

Quantity:	100 μL	
Target:	NOV	
Binding Specificity:	AA 139-357	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NOV antibody is un-conjugated	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)	
Product Details		
	NOV (Gly139-Met357)	
Immunogen:	NOV (Gly139-Met357)	
Immunogen: Isotype:	NOV (Gly139-Met357)	
Isotype:	IgG	
Isotype:	IgG  The antibody is a rabbit polyclonal antibody raised against NOV. It has been selected for its	
Isotype: Specificity:	IgG  The antibody is a rabbit polyclonal antibody raised against NOV. It has been selected for its ability to recognize NOV in immunohistochemical staining and western blotting.	
Isotype: Specificity: Purification:	IgG  The antibody is a rabbit polyclonal antibody raised against NOV. It has been selected for its ability to recognize NOV in immunohistochemical staining and western blotting.	
Isotype: Specificity: Purification: Target Details	IgG  The antibody is a rabbit polyclonal antibody raised against NOV. It has been selected for its ability to recognize NOV in immunohistochemical staining and western blotting.  Antigen-specific affinity chromatography	
Isotype: Specificity: Purification: Target Details Target:	IgG  The antibody is a rabbit polyclonal antibody raised against NOV. It has been selected for its ability to recognize NOV in immunohistochemical staining and western blotting.  Antigen-specific affinity chromatography	


	Protein NOV homolog	
Pathways:	Smooth Muscle Cell Migration, Growth Factor Binding	
Application Details		
Application Notes:	<ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500         Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.     </li> </ul>	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	

#### **Images**



### **Western Blotting**

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



#### Immunohistochemistry

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue