antibodies -online.com







Overview



Go to	D		
	Pron	HCL	nane

Quantity:	100 μL
Target:	NXF1
Binding Specificity:	AA 2-197
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NXF1 antibody is un-conjugated

Product Details

Application:

Immunogen:	NXF1 (Ala2-Tyr197)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against NXF1. It has been selected for its ability to recognize NXF1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Target Details

ranget Betano		
Target:	NXF1	
Abstract:	NXF1 Products	
Background:	Alternative Names: TAP, MEX67, Tip Associating Protein, Tip-associated protein, mRNA export	

factor TAP

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded liver tissue