antibodies - online.com







anti-OCEL1 antibody (AA 1-109)





| \sim | | | | |
|--------|------|-----|-----|-----|
| | 11/6 | ٦r١ | /10 | ۱۸. |

| Quantity: | 100 μL | |
|----------------------|--|--|
| Target: | OCEL1 | |
| Binding Specificity: | AA 1-109 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This OCEL1 antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) | |
| | | |

Product Details

| Immunogen: | OCEL1 (Met1-Ala109) | |
|---------------|--|--|
| Isotype: | IgG | |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against OCEL1. It has been selected for its ability to recognize OCEL1 in immunohistochemical staining and western blotting. | |
| Purification: | Antigen-specific affinity chromatography | |


Target Details

| Target: | OCEL1 | | |
|-------------------|---|--|--|
| Alternative Name: | Occludin/ELL Domain Containing Protein 1 (OCEL1) (OCEL1 Products) | | |

Application Details


| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. |
|--------------------|--|
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | Lot specific |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. |

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue