
antibodies -online.com





anti-OPA3 antibody (AA 1-200)

3 Images



Go to Product page

Ovarvian

Overview	
Quantity:	100 μL
Target:	OPA3
Binding Specificity:	AA 1-200
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OPA3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	OPA3 (Met1-Thr200)
Immunogen:	OPA3 (Met1-Thr200) IgG
Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against OPA3. It has been selected for its
Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against OPA3. It has been selected for its ability to recognize OPA3 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against OPA3. It has been selected for its ability to recognize OPA3 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against OPA3. It has been selected for its ability to recognize OPA3 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography

Aciduria Type 3

Application Details

	11	
Ann	lication	Notes.
, ,pp	Hoution	I VOLCO.

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images




Immunohistochemistry

Image 1. Used in DAB staining on fromalin fixed paraffinembedded kidney tissue



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. DAB staining on IHC-P; Samples: Rat Kidney Tissue