antibodies -online.com





Datasheet for ABIN5014065

anti-OS9 antibody (AA 471-672)





Go to Product page

\sim				
	$ V \cap$	r\/I	19	٨

Overview	
Quantity:	100 μL
Target:	OS9
Binding Specificity:	AA 471-672
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This OS9 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	

Immunogen:	OS9 (Asn471-Phe672)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against OS9. It has been selected for its ability to recognize OS9 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography


Target Details

Target:	OS9
Alternative Name:	Osteosarcoma Amplified 9 (OS9) (OS9 Products)
Background:	Alternative Names: ERLEC2, Amplified In Osteosarcoma, Endoplasmic Reticulum Lectin 2,

Target Details


nance of Protein Location, SARS-CoV-2 Protein Interactome tern blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 unohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in ffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal king dilutions must be determined by end user.
tern blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 unohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in ffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal
unohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in ffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal
unohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in ffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal
ermal stability is described by the loss rate. The loss rate was determined by accelerated all degradation test, that is, incubate the protein at 37°C for 48h, and no obvious ation and precipitation were observed. The loss rate is less than 5% within the expiration and appropriate storage condition.
search Use only
ecific

Images




Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. DAB staining on IHC-P; Samples: Mouse Testis Tissue



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded kidney tissue