# antibodies - online.com











_							
0	V	е	r١	/1	е	V	1

Overview	
Quantity:	100 μL
Target:	Pirin (PIR)
Binding Specificity:	AA 1-290
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Pirin antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	PIR (Met1-Asn290)

Immunogen:	PIR (Met1-Asn290)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against PIR. It has been selected for its ability to recognize PIR in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

# **Target Details**

Target:	Pirin (PIR)
Abstract:	PIR Products
Background:	Alternative Names: Iron Binding Nuclear Protein, Probable quercetin 2,3-dioxygenase PIR,

### Probable quercetinase

# **Application Details**

#### Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

#### Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

# Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

## **Images**



# **Western Blotting**

**Image 1.** Figure. Western Blot; Sample: Recombinant protein.



# **Immunohistochemistry**

**Image 2.** Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue