antibodies.com

# Datasheet for ABIN5014375 anti-SDCBP antibody (AA 2-298)

3 Images



#### Overview

| Quantity:            | 100 µL   |
|----------------------|--|
| Target:              | SDCBP  |
| Binding Specificity: | AA 2-298   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This SDCBP antibody is un-conjugated   |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) |

## Product Details

| Immunogen:    | ST1 (Ser2-Val298)  |
|---------------|--|
| Isotype:      | IgG  |
| Specificity:  | The antibody is a rabbit polyclonal antibody raised against ST1. It has been selected for its ability to recognize ST1 in immunohistochemical staining and western blotting. |
| Purification: | Antigen-specific affinity chromatography   |

#### Target Details


| Target:           | SDCBP   |
|-------------------|---|
| Alternative Name: | Syntenin 1 (ST1) (SDCBP Products)   |
| Background:       | Alternative Names: SDCBP, MDA9, SYCL, TACIP18, Syndecan Binding Protein, Scaffold protein |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5014375 | 09/10/2023 | Copyright antibodies-online. All rights reserved. Pbp1, Melanoma differentiation-associated protein 9, Pro-TGF-alpha cytoplasmic domaininteracting 18

## Application Details


| Application Notes: | <ul> <li>Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500<br/>Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in<br/>paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal<br/>working dilutions must be determined by end user.</li> </ul> |  |  |
|--------------------|---|--|--|
| Comment:           | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37&degC for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.        |  |  |
| Restrictions:      | For Research Use only   |  |  |
| Handling           |   |  |  |
| Format:            | Liquid  |  |  |
| Concentration:     | Lot specific  |  |  |
| Buffer:            | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.   |  |  |

#### Images



#### Western Blotting

**Image 1.** Western Blot; Sample: Human Urine; Primary Ab: 1µg/ml Rabbit Anti-Human ST1 Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



| Western Blotting |         |         |       |         |             |  |  |  |
|------------------|---------|---------|-------|---------|-------------|--|--|--|
| Image 2.         | Figure. | Western | Blot; | Sample: | Recombinant |  |  |  |
| protein.         |         |         |       |         |             |  |  |  |
|                  |         |         |       |         |             |  |  |  |

## Immunohistochemistry

**Image 3.** Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue

