antibodies -online.com







anti-TSTA3 antibody (AA 1-321)





\sim	
()\/\	rview
\cup	

Quantity:	100 μL
Target:	TSTA3
Binding Specificity:	AA 1-321
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TSTA3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	TSTA3 (Met1-Lys321)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TSTA3. It has been selected for its ability to recognize TSTA3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TSTA3
Abstract:	TSTA3 Products
Background:	Alternative Names: FX_P35B_SDR4F1_Short-chain dehydrogenase/reductase family 4F

member 1, GDP-L-Fucose Synthase, GDP-4-keto-6-deoxy-D-mannose-3,5-epimerase-4-reductase

Application Details

aaA	lication	Notes:
' 'PP	iioa tioi i	1.000.

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded kidney tissue