antibodies .- online.com







anti-TCN2 antibody (AA 19-427)





\sim				
	11/6	٦r١	/10	۱۸.

Quantity:	100 μL
Target:	TCN2
Binding Specificity:	AA 19-427
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TCN2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	TCN2 (Glu19-Met427)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TCN2. It has been selected for its ability to recognize TCN2 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TCN2
Alternative Name:	Transcobalamin II, Macrocytic Anemia (TCN2) (TCN2 Products)
Background:	Alternative Names: TC2. Transcobalamin II


Application Details

Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.
Images	

kDa 250-150-100-70-50-40-30-20-15-19-


Western Blotting

Image 1. Western Blot; Sample: Rat Serum; ;Primary Ab: 2μg/ml Rabbit Anti-Rat TCN2 Antibody;Second Ab: 0.2μg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody;(Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue