



[Go to Product page](#)

Datasheet for ABIN5014436
anti-TRERF1 antibody (AA 779-936)

2 Images

Overview

Quantity:	100 µL
Target:	TRERF1
Binding Specificity:	AA 779-936
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TRERF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	TRERF1 (Pro779-Arg936)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TRERF1. It has been selected for its ability to recognize TRERF1 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TRERF1
Abstract:	TRERF1 Products
Background:	Alternative Names: BCAR2, TREP132, RAPA, Breast Cancer Anti-Estrogen Resistance 2,

Target Details

Transcriptional-regulating protein 132, Zinc finger protein rapa, Zinc finger transcription factor TReP-132

Pathways:

[Regulation of Hormone Metabolic Process](#), [Regulation of Hormone Biosynthetic Process](#), [Chromatin Binding](#)

Application Details

Application Notes:

- Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only

Handling

Format:

Liquid


Concentration:

Lot specific

Buffer:


PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on formalin fixed paraffin-embedded esophagus tissue