antibodies -online.com







anti-TPM3 antibody (AA 6-248)

Images



()	11/0	r\ /1	$\triangle 1 $
	$\lor \lor \vdash$	1 V I	ew

Quantity:	100 μL
Target:	TPM3
Binding Specificity:	AA 6-248
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TPM3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)

Product Details

Immunogen:	TPM3 (Thr6-Met248)
Isotype:	IgG
Specificity:	The antibody is a rabbit polyclonal antibody raised against TPM3. It has been selected for its ability to recognize TPM3 in immunohistochemical staining and western blotting.
Purification:	Antigen-specific affinity chromatography

Target Details

Target:	TPM3	
Abstract:	TPM3 Products	
Background:	Alternative Names: NEM1, TRK, OK/SW-cl.5, TM-5, TM3, TM30, TM30nm, TPMsk3, Hscp30,	

Gamma-tropomyosin

Application Details

Application Notes:

• Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific

PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Concentration:


Images

Buffer:

kDa


Western Blotting

Image 1. Western Blot; Sample: Mouse Placenta lysate; Primary Ab: 2µg/ml Rabbit Anti-Mouse TPM3 Antibody Second Ab: 0.2µg/mL HRP-Linked Caprine Anti-Rabbit IgG Polyclonal Antibody (Catalog: SAA544Rb19)



Western Blotting

Image 2. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 3. Used in DAB staining on fromalin fixed paraffinembedded spleen tissue