

Datasheet for ABIN5014482

anti-DR3/LARD antibody (AA 38-167)

2 Images



Go to Product page


_				
()	ve.	rv/	101	Λ

Quantity:	100 μL
Target:	DR3/LARD (TNFRSF25)
Binding Specificity:	AA 38-167
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This DR3/LARD antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	TNFRSF25 (Cys38-Gly167)
Immunogen:	TNFRSF25 (Cys38-Gly167)
Isotype:	IgG
Isotype:	IgG The antibody is a rabbit polyclonal antibody raised against TNFRSF25. It has been selected for
Isotype: Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against TNFRSF25. It has been selected for its ability to recognize TNFRSF25 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification:	IgG The antibody is a rabbit polyclonal antibody raised against TNFRSF25. It has been selected for its ability to recognize TNFRSF25 in immunohistochemical staining and western blotting.
Isotype: Specificity: Purification: Target Details	IgG The antibody is a rabbit polyclonal antibody raised against TNFRSF25. It has been selected for its ability to recognize TNFRSF25 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography
Isotype: Specificity: Purification: Target Details Target:	IgG The antibody is a rabbit polyclonal antibody raised against TNFRSF25. It has been selected for its ability to recognize TNFRSF25 in immunohistochemical staining and western blotting. Antigen-specific affinity chromatography DR3/LARD (TNFRSF25)

Target Details


	Translocating Chain-Association Membrane Protein, Lymphocyte-associated receptor of death	
Pathways:	Apoptosis	
Application Details		
Application Notes:	 Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. 	
Comment:	The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	Lot specific	
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.	

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded intestine tissue