antibodies -online.com





anti-UMPS antibody (AA 205-480)






| | do to i roddot page |
|--|---------------------|
| | |
| | |

| Overview | | |
|----------------------|--|--|
| Quantity: | 100 μL | |
| Target: | UMPS | |
| Binding Specificity: | AA 205-480 | |
| Reactivity: | Human | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This UMPS antibody is un-conjugated | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC) | |
| Product Details | | |
| Immunogen: | UMPS (Val205-Val480) | |
| Isotype: | IgG | |
| Specificity: | The antibody is a rabbit polyclonal antibody raised against UMPS. It has been selected for its ability to recognize UMPS in immunohistochemical staining and western blotting. | |
| Purification: | Antigen-specific affinity chromatography | |
| Target Details | | |
| Target: | UMPS | |
| Abstract: | UMPS Products | |
| Background: | Alternative Names: OPRT, Orotate Phosphoribosyl Transferase, Orotidine-5'-Decarboxylase, | |


| | Orotate phosphoribosyltransferase, Orotidine 5'-phosphate decarboxylase | |
|---------------------|--|--|
| Pathways: | Ribonucleoside Biosynthetic Process | |
| Application Details | | |
| Application Notes: | Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user. | |
| Comment: | The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | Lot specific | |
| Buffer: | PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol. | |

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded Kidney tissue