
antibodies .- online.com





anti-UPP1 antibody (AA 45-235)

2 Images



Go to Product page

)\/(

Abstract:

Background:

Quantity:	100 μL
Target:	UPP1
Binding Specificity:	AA 45-235
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This UPP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC)
Product Details	
Immunogen:	UPP1 (Asn45-Tyr235)
la atoma a	
Isotype:	IgG
Specificity:	IgG The antibody is a rabbit polyclonal antibody raised against UPP1. It has been selected for its ability to recognize UPP1 in immunohistochemical staining and western blotting.
	The antibody is a rabbit polyclonal antibody raised against UPP1. It has been selected for its
Specificity:	The antibody is a rabbit polyclonal antibody raised against UPP1. It has been selected for its ability to recognize UPP1 in immunohistochemical staining and western blotting.
Specificity: Purification:	The antibody is a rabbit polyclonal antibody raised against UPP1. It has been selected for its ability to recognize UPP1 in immunohistochemical staining and western blotting.

Alternative Names: UDRPASE, UP, UPASE

UPP1 Products

Pathways:

Ribonucleoside Biosynthetic Process

Application Details

Application Notes:

Western blotting: 1:50-400 Immunocytochemistry in formalin fixed cells: 1:50-500
 Immunohistochemistry in formalin fixed frozen section: 1:50-500 Immunohistochemistry in paraffin section: 1:10-100 Enzyme-linked Immunosorbent Assay: 1:100-1:5000 Optimal working dilutions must be determined by end user.

Comment:

The thermal stability is described by the loss rate. The loss rate was determined by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.


Restrictions:

For Research Use only

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	PBS, pH 7.4, containing 0.02 % Sodium azide, 50 % glycerol.

Images



Western Blotting

Image 1. Figure. Western Blot; Sample: Recombinant protein.



Immunohistochemistry

Image 2. Used in DAB staining on fromalin fixed paraffinembedded pancreas tissue