

Datasheet for ABIN502122 anti-CKMT2 antibody (C-Term)

2 Images



Go to Product page

\sim				
()\	10	rVI	161	٨
\cup	ノロ	V	\Box	Λ

Quantity:	25 μL
Target:	CKMT2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Rabbit, Cow, Horse, Zebrafish (Danio rerio), Guinea Pig, Saccharomyces cerevisiae
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This CKMT2 antibody is un-conjugated
Application:	Western Blotting (WB)
Product Details	
Immunogen:	The immunogen is a synthetic peptide directed towards the C terminal region of human CKMT2
Sequence:	ISNIDRIGRS EVELVQIVID GVNYLVDCEK KLERGQDIKV PPPLPQFGKK
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 93%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%, Yeast: 80%, Zebrafish: 83%
Characteristics:	This is a rabbit polyclonal antibody against CKMT2. It was validated on Western Blot using a cell lysate as a positive control.
Purification:	Affinity Purified
Target Details	
Target:	CKMT2

Alternative Name:	CKMT2 (CKMT2 Products)
Background:	Mitochondrial creatine kinase (MtCK) is responsible for the transfer of high energy phosphate
	from mitochondria to the cytosolic carrier, creatine. It belongs to the creatine kinase isoenzym
	family. It exists as two isoenzymes, sarcomeric MtCK and ubiquitous MtCK, encoded by
	separate genes. Mitochondrial creatine kinase occurs in two different oligomeric forms: dimer
	and octamers, in contrast to the exclusively dimeric cytosolic creatine kinase isoenzymes.
	Sarcomeric mitochondrial creatine kinase has 80 % homology with the coding exons of
	ubiquitous mitochondrial creatine kinase. This gene contains sequences homologous to
	several motifs that are shared among some nuclear genes encoding mitochondrial proteins
	and thus may be essential for the coordinated activation of these genes during mitochondrial
	biogenesis. Three transcript variants encoding the same protein have been found for this
	gene.Mitochondrial creatine kinase (MtCK) is responsible for the transfer of high energy
	phosphate from mitochondria to the cytosolic carrier, creatine. It belongs to the creatine kinas
	isoenzyme family. It exists as two isoenzymes, sarcomeric MtCK and ubiquitous MtCK,
	encoded by separate genes. Mitochondrial creatine kinase occurs in two different oligomeric
	forms: dimers and octamers, in contrast to the exclusively dimeric cytosolic creatine kinase
	isoenzymes. Sarcomeric mitochondrial creatine kinase has 80 % homology with the coding
	exons of ubiquitous mitochondrial creatine kinase. This gene contains sequences homologous
	to several motifs that are shared among some nuclear genes encoding mitochondrial proteins
	and thus may be essential for the coordinated activation of these genes during mitochondrial
	biogenesis. Three transcript variants encoding the same protein have been found for this gene
	Alias Symbols: SMTCK
	Protein Interaction Partner: UBC, ABHD6, TMED9, ELN, PSMD4, LRIF1, OLFML3, UNC119,
	CKMT2,
	Protein Size: 419
Molecular Weight:	43 kDa
Gene ID:	1160
NCBI Accession:	NM_001825, NP_001816
UniProt:	P17540
Application Details	
Application Notes:	Optimal working dilutions should be determined experimentally by the investigator.
Comment:	Antigen size: 419 AA


Application Details

Restrictions:	For Research Use only
---------------	-----------------------

Handling


Format:	Liquid
Concentration:	Lot specific
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Avoid repeated freeze-thaw cycles.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.

Images



Western Blotting

Image 1. Host: Rabbit Target Name: CKMT2 Sample Type: Human Fetal Muscle Antibody Dilution: 1.0ug/ml



Western Blotting

Image 2. WB Suggested Anti-CKMT2 Antibody Titration: 0.2-1 ug/ml ELISA Titer: 1:312500 Positive Control: Human heart