

Datasheet for ABIN5066211  
**anti-ATG9B antibody (AA 110-121) (Atto 488)**[Go to Product page](#)

## 2 Images

## Overview

Quantity:	100 µg
Target:	ATG9B
Binding Specificity:	AA 110-121
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ATG9B antibody is conjugated to Atto 488
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)

## Product Details

Immunogen:	Synthetic peptide from the N-terminal of Human ATG9B (aa. 110-121)
Isotype:	IgG
Specificity:	Highly expressed in placenta (trophoblast cells) and pituitary gland. Not expressed in vascular endothelial., Detects ~100 kDa.
Cross-Reactivity:	Human, Mouse
Purification:	Peptide Affinity Purified

## Target Details

Target:	ATG9B
Alternative Name:	ATG9B ( <a href="#">ATG9B Products</a> )

## Target Details


Background:	ATG9B is involved in autophagy and cytoplasm to vacuole transport (Cvt) vesicle formation. Plays a key role in the organization of the preautophagosomal structure/phagophore assembly site (PAS), the nucleating site for formation of the sequestering vesicle. Highly expressed in the placenta and the pituitary gland.
Gene ID:	285973
NCBI Accession:	<a href="#">NP_001303985</a>
UniProt:	<a href="#">Q674R7</a>

## Application Details

Application Notes:	<ul style="list-style-type: none"><li>• WB (1:1000)</li><li>• ICC/IF (1:100)</li><li>• optimal dilutions for assays should be determined by the user.</li></ul>
Comment:	A 1:1000 dilution of ABIN5066211 was sufficient for detection of ATG9B in 15 µg of Human HeLa Cell Lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only


## Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C



#### Immunofluorescence (fixed cells)

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-ATG9B Polyclonal Antibody . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-ATG9B Polyclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Cytoplasmic Vesicle, Autophagosome Membrane, Multi-Pass Membrane Protein. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) ATG9B Antibody (D) Composite.



#### Western Blotting

**Image 2.** Western blot analysis of Rat Brain cell lysates showing detection of 101 kDa ATG9B protein using Rabbit Anti-ATG9B Polyclonal Antibody . Lane 1: Molecular Weight Ladder (MW). Lane 2: Rat Brain cell lysates. Load: 15 µg . Primary Antibody: Rabbit Anti-ATG9B Polyclonal Antibody at 1:1000 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: 101 kDa.