

Datasheet for ABIN5066674  
**anti-KCNJ10 antibody (PerCP)**[Go to Product page](#)

## 3 Images

## Overview

|              |   |
|--------------|---|
| Quantity:    | 100 µg  |
| Target:      | KCNJ10  |
| Reactivity:  | Human   |
| Host:        | Rabbit  |
| Clonality:   | Polyclonal  |
| Conjugate:   | This KCNJ10 antibody is conjugated to PerCP   |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF) |

## Product Details

|                   |  |
|-------------------|--|
| Immunogen:        | Synthetic peptide from the mid-protein of Human Kir4.1   |
| Isotype:          | IgG  |
| Specificity:      | Expressed in kidney (at protein level), Detects ~42 kDa. |
| Cross-Reactivity: | Human, Rat   |
| Purification:     | Peptide Affinity Purified                                |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | KCNJ10                                     |
| Alternative Name: | Kir4.1 ( <a href="#">KCNJ10 Products</a> ) |
| Gene ID:          | 3766                                       |

## Target Details

|                 |  |
|-----------------|--|
| NCBI Accession: | <a href="#">NP_002232</a>  |
| UniProt:        | <a href="#">P78508</a>   |
| Pathways:       | <a href="#">Dicarboxylic Acid Transport</a> , <a href="#">Regulation of long-term Neuronal Synaptic Plasticity</a> |

## Application Details

|                    |   |
|--------------------|---|
| Application Notes: | <ul style="list-style-type: none"><li>• WB (1:1000)</li><li>• ICC/IF (1:100)</li><li>• IHC (1:50)</li><li>• optimal dilutions for assays should be determined by the user.</li></ul>            |
| Comment:           | A 1:1000 dilution of ABIN5066674 was sufficient for detection of Kir4.1 in 15 µg of rat liver cell lysates by ECL immunoblot analysis using goat anti-rabbit IgG:HRP as the secondary antibody. |
| Restrictions:      | For Research Use only   |


## Handling

|                    |  |
|--------------------|--|
| Format:            | Liquid   |
| Concentration:     | 1 mg/mL  |
| Buffer:            | PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated                                     |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | 4 °C   |
| Storage Comment:   | Conjugated antibodies should be stored at 4°C  |




### Immunohistochemistry

**Image 1.** Immunohistochemistry analysis using Rabbit Anti-Kir4.1 Polyclonal Antibody (ABIN5066674). Tissue: Kidney. Species: Human. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-Kir4.1 Polyclonal Antibody (ABIN5066674) at 1:50 for 30 min at RT. Counterstain: Hematoxylin. Magnification: 10X. HRP-DAB Detection.



### Western Blotting

**Image 2.** Western blot analysis of Rat Liver cell lysates showing detection of ~42.5 kDa Kir4.1 protein using Rabbit Anti-Kir4.1 Polyclonal Antibody. Lane 1: Molecular Weight Ladder (MW). Lane 2: Rat Liver cell lysates. Load: 15 µg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Rabbit Anti-Kir4.1 Polyclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Rabbit IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~42.5 kDa.



### Immunofluorescence (fixed cells)

**Image 3.** Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Kir4.1 Polyclonal Antibody. Tissue: Colon carcinoma cell line (RKO). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Rabbit Anti-Kir4.1 Polyclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Rabbit ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Membrane, Cytoplasm. Magnification: 60X. (A) DAPI nuclear stain. (B) Phalloidin Texas Red F-Actin stain. (C) Kir4.1 Antibody. (D) Composite.