

Datasheet for ABIN5066877

**anti-SCNN1B antibody (AA 617-638) (Atto 488)**[Go to Product page](#)**1** Image

## Overview

Quantity:	100 µg
Target:	SCNN1B
Binding Specificity:	AA 617-638
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SCNN1B antibody is conjugated to Atto 488
Application:	Western Blotting (WB), Immunohistochemistry (IHC)

## Product Details

Immunogen:	Synthetic peptide from the C-terminal of Rat ENaC beta (aa. 617-638)
Clone:	7B8
Isotype:	IgG1
Specificity:	Detects ~87 kDa.
Cross-Reactivity:	Mouse
Purification:	Protein G Purified

## Target Details

Target:	SCNN1B
Alternative Name:	ENaC beta ( <a href="#">SCNN1B Products</a> )

## Target Details

**Background:** The Epithelial Sodium Channel (ENaC) is a membrane ion channel permeable to Na<sup>+</sup> ions. It is located in the apical plasma membrane of epithelia in the kidneys, lung, colon, and other tissues where it plays a role in trans epithelial Na<sup>+</sup>-ion transport (1). Specifically Na<sup>+</sup> transport via ENaC occurs across many epithelial surfaces, and plays a key role in regulating salt and water absorption (2). ENaCs are composed of three structurally related subunits that form a tetrameric channel, alpha, beta, and gamma. The expression of its alpha and beta subunits is enhanced as keratinocytes differentiate (3, 4). The beta and gamma-ENaC subunits are essential for edema fluid to exert its maximal effect on net fluid absorption by distal lung epithelia(5). And it has been concluded that the subunits are differentially expressed in the retina of mice with ocular hypertension, therefore the up-regulation of alpha-ENaC proteins could serve as a protection mechanism against elevated intraocular pressure (6).

**Gene ID:** 24767

**NCBI Accession:** [NP\\_036780](#)

**UniProt:** [P37090](#)

## Application Details

**Application Notes:**

- WB (1:1000)
- IHC (1:150)
- optimal dilutions for assays should be determined by the user.

**Comment:** A 1:1000 dilution of ABIN5066877 was sufficient for detection of ENaC beta in 15 µg of Mouse whole kidney lysate by ECL immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

**Restrictions:** For Research Use only

## Handling

**Format:** Liquid

**Concentration:** 1 mg/mL

**Buffer:** PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

**Preservative:** Sodium azide


**Precaution of Use:** This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

## Handling

Storage: 4 °C

Storage Comment: Conjugated antibodies should be stored at 4°C

## Images



### Immunohistochemistry

**Image 1.** Immunohistochemistry analysis using Mouse Anti-ENaC beta Monoclonal Antibody, Clone 7B8 . Tissue: Kidney (cortex). Species: Mouse. Primary Antibody: Mouse Anti-ENaC beta Monoclonal Antibody at 1:150. Localization: Collecting duct principal cells. Magnification: 60X.