



[Go to Product page](#)

Datasheet for ABIN5067356

## anti-Malondialdehyde antibody (HRP)

### 3 Images

#### Overview

|              |   |
|--------------|---|
| Quantity:    | 100 µg  |
| Target:      | Malondialdehyde (MDA)   |
| Reactivity:  | Please inquire  |
| Host:        | Mouse   |
| Clonality:   | Monoclonal  |
| Conjugate:   | This Malondialdehyde antibody is conjugated to HRP  |
| Application: | Western Blotting (WB), ELISA, Immunocytochemistry (ICC), Flow Cytometry (FACS), Immunofluorescence (IF) |

#### Product Details

|               |  |
|---------------|--|
| Immunogen:    | Synthetic Malondialdehyde modified Keyhole Limpet Hemocyanin (KLH).  |
| Clone:        | 6H6  |
| Isotype:      | IgG1   |
| Specificity:  | Specific for MDA conjugated proteins. Does not detect free MDA. Does not cross-react with Acrolein, Crotonaldehyde, Hexanoyl Lysine, 4-HHE, 4-HNE, or Methylglyoxal modified proteins. |
| Purification: | Protein G Purified   |

#### Target Details

|                   |  |
|-------------------|--|
| Target:           | Malondialdehyde (MDA)                            |
| Alternative Name: | Malondialdehyde ( <a href="#">MDA Products</a> ) |

## Target Details

---

Target Type: Chemical

---

Background: Malondialdehyde (MDA) is the biomarker in greatest diagnostic use, due to its molecular stability. This three-carbon, low-molecular weight aldehyde has a strong affinity for amino acids, which results in adduct formation to both free amino acids and proteins. Increased MDA levels have been found at correlating levels in breast cancer, and lung cancer patients. Other diseased states with elevated MDA levels include diabetes and Alzheimer's disease. Multiple laboratory techniques exist for quantification of MDA levels, including the thiobarbituric acid reactive substances (TBARS) assay. In addition to use as a biomarker, MDA has been shown to have mutagenic effects on tissues themselves as adduct formation can result in DNA cross-linking.

## Application Details

---

Application Notes:

- WB (1:1000)
- ICC/IF (1:50)
- FACS (1:50)
- FCM (1:50)
- ELISA (1:1000)
- optimal dilutions for assays should be determined by the user.

---

Comment: A 1:1000 dilution of ABIN5067356 was sufficient for detection of Malondialdehyde in 2 µg of Malondialdehyde conjugated to BSA by ECL immunoblot analysis using Goat Anti-Mouse IgG:HRP as the secondary Antibody.

---

Restrictions: For Research Use only

---

## Handling

---

Format: Liquid

---

Concentration: 1 mg/mL

---

Buffer: PBS pH 7.4, 50 % glycerol, 0.09 % Sodium azide, Storage buffer may change when conjugated

---

Preservative: Sodium azide

---

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


---

Storage: 4 °C

---


Storage Comment: Conjugated antibodies should be stored at 4°C

---




### Immunofluorescence (fixed cells)

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6 . Tissue: Embryonic kidney cells (HEK293). Species: Human. Fixation: 5% Formaldehyde for 5 min. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:50 for 30-60 min at RT. Secondary Antibody: Goat Anti-Mouse Alexa Fluor 488 at 1:1500 for 30-60 min at RT. Counterstain: Phalloidin Alexa Fluor 633 F-Actin stain; DAPI (blue) nuclear stain at 1:250, 1:50000 for 30-60 min at RT. Magnification: 20X (2X Zoom). (A,C,E,G) - Untreated. (B,D,F,H) - Cells cultured overnight with 50  $\mu$ M H<sub>2</sub>O<sub>2</sub>. (A,B) DAPI (blue) nuclear stain. (C,D) Phalloidin Alex Fluor 633 F-Actin stain. (E,F) Malondialdehyde Antibody. (G,H) Composite. Courtesy of: Dr. Robert Burke, University of Victoria.



### Western Blotting

**Image 2.** Western Blot analysis of Malondialdehyde-BSA Conjugate showing detection of 67 kDa Malondialdehyde - BSA using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6 . Lane 1: Molecular Weight Ladder (MW). Lane 2: Malondialdehyde-BSA (0.5  $\mu$ g). Lane 3: Malondialdehyde-BSA (2.0  $\mu$ g). Lane 4: BSA (0.5  $\mu$ g). Lane 5: BSA (2.0  $\mu$ g) . Block: 5% Skim Milk in TBST. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 67 kDa.



### Flow Cytometry

**Image 3.** Flow Cytometry analysis using Mouse Anti-Malondialdehyde Monoclonal Antibody, Clone 6H6. Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 90% Methanol. Primary Antibody: Mouse Anti-Malondialdehyde Monoclonal Antibody at 1:50 for 30 min on ice. Secondary Antibody: Goat Anti-Mouse: PE at 1:100 for 20 min at RT. Cells were subject to oxidative stress by treating with 250  $\mu$ M H<sub>2</sub>O<sub>2</sub> for 24 hours.