

Datasheet for ABIN5067873

anti-CD86 antibody





Overview

Quantity:	100 μg
Target:	CD86
Reactivity:	Human, Cynomolgus, Baboon, Rhesus Monkey
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CD86 antibody is un-conjugated
Application:	Flow Cytometry (FACS)

Product Details

Purpose:

Immunogen:

Clone:	BU63
Isotype:	IgG1, kappa
Characteristics:	BU63 recognizes ~70Kda a surface antigen protein of Ig superfamily characterized as CD86
	expressed by most antigen presenting cells (APC). Depending on the immunological condition,
	CD86 specifically binds with its two ligands CD28 and CD152 (CTLA-4) expressed by T cells.
	Binding with CD28, CD86 provides stimulatory signal and activates T cells whereas binding with
	CD152 (CTLA-4) CD86 provides inhibitory signal to T cells. Generally, high-level CD86
	expression was noticed on peripheral monocytes, tissue macrophages and dendritic cells, and
	also on activated B cells. CD86 is useful in the clinical diagnosis of cancer cells including
	malignant Hodgkin's and non-hodgkin's diseases.

CD86 Unconjugated Antibody

B-lymphoblastoid cell line ARH 77

Product Details

Purification:	Purified
Purity:	>95 %
Grade:	GMP Grade

Target Details


Target:	CD86
Alternative Name:	CD86 (CD86 Products)
UniProt:	P42081
Pathways:	TCR Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Activation of Innate immune Response, Cellular Response to Molecule of
	Bacterial Origin, Positive Regulation of Immune Effector Process, Activated T Cell Proliferation

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Buffer:	PBS pH 7.2, 0.1 % (w/v) BSA, 0.09 % (w/v) sodium azide
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C



Flow Cytometry

Image 1.