antibodies.com

Datasheet for ABIN513555 anti-Aspartate beta Hydroxylase antibody (AA 1-210)



1 Image

Overview

Quantity:	50 µg
Target:	Aspartate beta Hydroxylase (ASPH)
Binding Specificity:	AA 1-210
Reactivity:	Human
Host:	Mouse
Clonality:	Polyclonal
Conjugate:	This Aspartate beta Hydroxylase antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details


Purpose:	Mouse polyclonal antibody raised against a full-length human ASPH protein.
Immunogen:	ASPH (NP_115856.1, 1 a.a. ~ 210 a.a) full-length human protein.
Sequence:	MAEDKETKHG GHKNGRKGGL SGTSFFTWFM VIALLGVWTS VAVVWFDLVD YEEVLGKLGI YDADGDGDFD VDDAKVLLEG PSGVAKRKTK AKVKELTKEE LKKEKEKPES RKESKNEERK KGKKEDVRKD KKIADADLSR KESPKGKKDR EKEKVDLEKS AKTKENRKKS TNMKDVSSKM ASRDKDDRKE SRSSTRYAHL TKGNTQKRNG
Cross-Reactivity:	Human
Characteristics:	Antibody reactive against mammalian transfected lysate.
Target Details	
Target:	Aspartate beta Hydroxylase (ASPH)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN513555 | 02/07/2024 | Copyright antibodies-online. All rights reserved.

Target Details		
Alternative Name:	ASPH (ASPH Products)	
Background:	Full Gene Name: aspartate beta-hydroxylase	
	Synonyms: BAH,CASQ2BP1,HAAH,JCTN,junctin	
Gene ID:	444	
NCBI Accession:	NM_032467	
Pathways:	Positive Regulation of Endopeptidase Activity	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Buffer:	In 1x PBS, pH 7.4	
Handling Advice:	Aliquot to avoid repeated freezing and thawing.	
Storage:	-20 °C	

Images

Storage Comment:



Western Blotting

Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Image 1. Western Blot analysis of ASPH expression in transfected 293T cell line by ASPH MaxPab polyclonal antibody.

Lane 1: ASPH transfected lysate(23.1 KDa). Lane 2: Non-transfected lysate.