

Datasheet for ABIN531693  
**anti-TICAM2 antibody (AA 1-235)**



[Go to Product page](#)

2 Images

## Overview

Quantity:	100 µL
Target:	TICAM2
Binding Specificity:	AA 1-235
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TICAM2 antibody is un-conjugated
Application:	Western Blotting (WB), Immunoprecipitation (IP)

## Product Details

Purpose:	Rabbit polyclonal antibody raised against a full-length human TICAM2 protein.
Immunogen:	TICAM2 (NP_067681.1, 1 a.a. ~ 235 a.a) full-length human protein.
Sequence:	MGIGKSKINS CPLSLSWGKR HSDVTSPGYH EDSKKS EDL SLCNVAEHSN TTEGPTGKQE GAQSVEEMFE EEAEVEFLK FVILHAEDDT DEALRVQNL QDDFGIKPGI IFAEMPCGRQ HLQNLDDAVN GSAWTILLT ENFLRDTWCN FQFYTSLMNS VNRQHKYNSV IPMRPLNNPL PRERTPFALQ TINAL EEE SR GFPTQVERIF QESVYKTQQT IWKETRNMVQ RQFIA
Cross-Reactivity:	Human
Characteristics:	Antibody reactive against mammalian transfected lysate.

## Target Details

Target:	TICAM2
---------	--------

## Target Details

Alternative Name:	TICAM2 ( <a href="#">TICAM2 Products</a> )
Background:	Full Gene Name: toll-like receptor adaptor molecule 2 Synonyms: MGC129876,MGC129877,TICAM-2,TIRAP3,TIRP,TRAM
Gene ID:	353376
NCBI Accession:	<a href="#">NM_021649</a>
Pathways:	<a href="#">TLR Signaling</a> , <a href="#">Activation of Innate immune Response</a> , <a href="#">Cellular Response to Molecule of Bacterial Origin</a> , <a href="#">Toll-Like Receptors Cascades</a>


## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	No additive
Preservative:	Without preservative
Handling Advice:	Aliquot to avoid repeated freezing and thawing.
Storage:	-20 °C
Storage Comment:	Store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

## Images




### Western Blotting

**Image 1.** Western Blot analysis of TICAM2 expression in transfected 293T cell line by TICAM2 MaxPab polyclonal antibody.

Lane 1: TICAM2 transfected lysate(26.90 KDa).

Lane 2: Non-transfected lysate.



### Immunoprecipitation

**Image 2.** Immunoprecipitation of TICAM2 transfected lysate using anti-TICAM2 MaxPab rabbit polyclonal antibody and Protein A Magnetic Bead , and immunoblotted with TICAM2 purified MaxPab mouse polyclonal antibody (B01P)