antibodies -online.com





anti-MAP2K5 antibody (C-Term)





Publication



Go to Product page

Overview

Quantity:	100 μg
Target:	MAP2K5
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat, Dog, Chicken
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP2K5 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunoprecipitation (IP)

Product Details

Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of MAP2K5.
Immunogen:	A synthetic peptide corresponding to C-terminus of human MAP2K5.
Specificity:	This antibody recognizes ~49 KDa of human MAP2K5.
Cross-Reactivity:	Chicken, Dog, Human, Mouse, Rat
Characteristics:	Antibody Reactive Against Synthetic Peptide.

Target Details

Target:	MAP2K5
Alternative Name:	MAP2K5 (MAP2K5 Products)
Gene ID:	5607

Target Details Pathways:

MAPK Signaling, Neurotrophin Signaling Pathway

Application Details

Application Notes:

Western Blot (0.1-1 µg/mL)

ELISA (0.01-0.1 μg/mL)

Immunoprecipitation (2-5 µg/mL)
Immunohistochemistry (2-5 µg/mL)

The optimal working dilution should be determined by the end user.

Restrictions:

For Research Use only


Handling

Format:	Liquid
Buffer:	In TBS, pH 7.2 (BSA, 10 % Proclin300)
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C. For long term storage store at -20°C or lower. Aliquot to avoid repeated freezing and thawing.

Publications

Product cited in:

Shishido, Woo, Ding, McClain, Molina, Yan, Yang, Abe: "Effects of MEK5/ERK5 association on small ubiquitin-related modification of ERK5: implications for diabetic ventricular dysfunction after myocardial infarction." in: **Circulation research**, Vol. 102, Issue 11, pp. 1416-25, (2008) (PubMed).



Western Blotting

Image 1. The tissue lysates derived from mouse liver were immunoblotted by MAP2K5 polyclonal antibody at 1 : 1000.