# antibodies -online.com





Datasheet for ABIN544032

# anti-EVX2 antibody (C-Term)

1 Image

3

**Publications** 



Go to Product page

#### Overview

Quantity:	400 μL
Target:	EVX2
Binding Specificity:	C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EVX2 antibody is un-conjugated
Application:	Western Blotting (WB)


### **Product Details**

Purpose:	Rabbit polyclonal antibody raised against synthetic peptide of Evx2.
Immunogen:	A synthetic peptide (conjugated with KLH) corresponding to C-terminus of mouse Evx2.
Cross-Reactivity:	Human, Mouse
Target Details	
Target:	EVX2

Target:	EVX2
Alternative Name:	EVX2 (EVX2 Products)
Gene ID:	14029

## **Application Details**

Application Notes:	Western Blot (1:1000)
	The optimal working dilution should be determined by the end user.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Buffer:	In PBS (0.09 % sodium azide)
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4°C. For long term storage store at -20°C.
	Aliquot to avoid repeated freezing and thawing.
Publications	
Product cited in:	Montavon, Le Garrec, Kerszberg, Duboule: "Modeling Hox gene regulation in digits: reverse
	collinearity and the molecular origin of thumbness." in: Genes & development, Vol. 22, Issue 3,
	pp. 346-59, (2008) (PubMed).
	Tarchini, Duboule: "Control of Hoxd genes' collinearity during early limb development." in:
	<b>Developmental cell</b> , Vol. 10, Issue 1, pp. 93-103, (2006) (PubMed).
	Tarchini, Huynh, Cox, Duboule: "HoxD cluster scanning deletions identify multiple defects
	leading to paralysis in the mouse mutant Ironside." in: <b>Genes &amp; development</b> , Vol. 19, Issue 23,
	pp. 2862-76, (2005) (PubMed).



#### **Western Blotting**

**Image 1.** Western blot analysis of Evx2 polyclonal antibody in mouse kidney tissue lysates (35 ug/lane). Evx2 (arrow) was detected using the purified polyclonal antibody.