

Datasheet for ABIN5508784
anti-beta Amyloid antibody (C-Term)



[Go to Product page](#)

2 Validations

4 Images

4 Publications

Overview

Quantity:	100 µg
Target:	beta Amyloid (Abeta)
Binding Specificity:	C-Term
Reactivity:	Human, Mouse, Rat
Host:	Mouse
Clonality:	Monoclonal
Application:	Western Blotting (WB), ELISA, Immunofluorescence (IF)

Product Details

Immunogen:	Synthetic peptide derived from Human Abeta 1-42 protein
Clone:	9E8-1A3
Isotype:	IgG1
Specificity:	A-beta 1-40, A-beta 1-42, A-beta 1-43 and APP
Cross-Reactivity (Details):	Human, mouse, rat (other species not tested)
Purification:	Protein G affinity purification

Target Details

Target:	beta Amyloid (Abeta)
Alternative Name:	Amyloid beta (Abeta Products)
Background:	Amyloid beta (Aβ or Abeta) is a peptide of 36 - 43 amino acids derived from the cleavage of

Target Details

APP (Amyloid Precursor Protein) by β - and γ -secretases. It is involved in Alzheimer's disease, among other severe neurological disorders. It has been proved that the A β could be a crucial target for these important diseases treatment.

Pathways: [Inflammasome](#)

Application Details

Application Notes: ELISA: 1:40,000, WB: 1:4,000, IF: 1:2,000

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS, pH 7.4

Preservative: Without preservative

Handling Advice: Avoid freezing/thaw cycles.

Storage: -20 °C

Storage Comment: Store at -20°C to -80°C.

Publications

Product cited in: Ordóñez-Gutiérrez, Posado-Fernández, Ahmadvand, Lettieri, Wu, Antón, Flores, Moghimi, Wandosell: "ImmunoPEGLiposome-mediated reduction of blood and brain amyloid levels in a mouse model of Alzheimer's disease is restricted to aged animals." in: **Biomaterials**, Vol. 112, pp. 141-152, (2016) ([PubMed](#)).


Markoutsa, Papadia, Clemente, Flores, Antimisiaris: "Anti-A β -MAb and dually decorated nanoliposomes: effect of A β 1-42 peptides on interaction with hCMEC/D3 cells." in: **European journal of pharmaceutics and biopharmaceutics : official journal of Arbeitsgemeinschaft für Pharmazeutische Verfahrenstechnik e.V.**, Vol. 81, Issue 1, pp. 49-56, (2012) ([PubMed](#)).

Le Droumaguet, Nicolas, Brambilla, Mura, Maksimenko, De Kimpe, Salvati, Zona, Airoldi, Canovi, Gobbi, Magali, La Ferla, Nicotra, Scheper, Flores, Masserini, Andrieux, Couvreur: "Versatile and efficient targeting using a single nanoparticulate platform: application to cancer and

Alzheimer's disease." in: **ACS nano**, Vol. 6, Issue 7, pp. 5866-79, (2012) ([PubMed](#)).


Canovi, Markoutsas, Lazar, Pampalakis, Clemente, Re, Sesana, Masserini, Salmona, Duyckaerts, Flores, Gobbi, Antimisiaris: "The binding affinity of anti-A β 1-42 MAb-decorated nanoliposomes to A β 1-42 peptides in vitro and to amyloid deposits in post-mortem tissue." in: **Biomaterials**, Vol. 32, Issue 23, pp. 5489-97, (2011) ([PubMed](#)).

Images




Western Blotting

Image 1. Western blots of the monoclonal antibody binding to different amyloid beta (Abeta) regions of human and mouse protein, using 12% A/B (acrylamide/bisacrylamide) or 10% Tricine matrix. Lanes (1) human Abeta 1-40; (2) human Abeta1-42; (3) human Abeta1-43; (4) mouse Abeta1-40; (5) mouse Abeta1-42.



Immunofluorescence

Image 2. Fluorescent microscopy image (apoptome) of 6 months-old APP/PS1 mouse hippocampus and somatosensory cortex. Double staining: in blue, nuclei DAPI staining. In red, Amyloid plaques.



Immunofluorescence

Image 3. Immunofluorescence slices with the anti-amyloid beta antibody from different mouse genotypes. Wild type (WT), APP/PS1 and APP KO. In red, amyloid plaques.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN5508784.



Successfully validated (Immunohistochemistry (IHC))

by Prof. Merighi, Laboratory of Neurobiology, Department of Veterinary Sciences, University of Turin

Report Number: 104425


Date: May 12 2022

Target:	beta Amyloid
Lot Number:	190522A
Method validated:	Immunohistochemistry (IHC)
Positive Control:	Adult (24 months) mouse brain 3-month-old Taconic mouse Model 1349 brain fixed in 4% paraformaldehyde
Negative Control:	We incubated slices overnight with the blocking solution only and then processed them with the secondary antibody.
Notes:	The beta amyloid antibody (Amyloid beta) (C-Term) ABIN5508784 works in IHC-P, especially at higher concentrations (1:200 and 1:500), and without the use of any antigen retrieval treatment.
Primary Antibody:	ABIN5508784
Secondary Antibody:	goat anti-rabbit AF488-conjugated antibody (Invitrogen by Thermo Fisher Scientific, A11034, lot 1971418)
Protocol:	<ul style="list-style-type: none">• Perfuse mice with paraformaldehyde 4% in 0.1 M phosphate buffer pH 7.4 and post-fix in the same fixative for an additional 2 h at RT.• Wash, dehydrate, and embed samples in paraffin wax.• Wash several times with 0.01 M PBS.• Cut intestines and brain with a microtome into 6µm sections and mount on glass slides.• After paraffin removal, incubate sections for 1 h at RT in PBS containing 1% albumin from chicken egg white (Sigma, A5378) and 0.3% Triton-X-100 (BioRad, 161-0407, lot 00583) to block non-specific binding sites.• Incubate sections with primary anti-beta amyloid (C-Term) (antibodies Online, ABIN5508784, lot 190522A) diluted 1:200, 1:500, 1:1000, and 1:2000 in PBS-BSA-PLL ON at RT.• Wash sections 3x 5 min with 0.01 M PBS.• Incubate sections with secondary goat anti-rabbit AF488-conjugated antibody (Invitrogen by Thermo Fisher Scientific, A11034, lot 1971418) diluted 1:500 in 0.1 M PBS for 1 h at RT.• Wash sections 3x 5 min with 0.01M PBS.• Mount specimens in Fluoroshield (Sigma, F6182, lot MKCB0153V).• Acquire images with a fluorescence microscope and appropriate filter settings for AF488, e.g. Leica DM 6000B fluorescence microscope equipped with a digital camera at 40x

magnification.

Experimental Notes: Antigen retrieval treatment was also tested. In this case, sections were processed for microwave antigen retrieval for 10 min(95-100 °C) in 10 mM sodium citrate buffer (pH 6.0). After 20 min of spontaneous cooling, sections were washed twice for 5 min with distilled water and for 5 min with PBS.

Image for Validation report #104425



Validation image no. 1 for anti-Amyloid beta (Abeta) (C-Term) antibody (ABIN5508784)

Staining of beta-amyloid positive cells in the adult mouse brain using ABIN5508784 at 40x magnification.



Successfully validated (Immunohistochemistry (IHC))

by [Prof. Merighi](#), Laboratory of Neurobiology, Department of Veterinary Sciences, University of Turin


Report Number: 104493

Date: Mar 15 2023

Target:	Abeta
Lot Number:	190522A
Method validated:	Immunohistochemistry (IHC)
Positive Control:	3-month-old Taconic mouse Model 1349 brain fixed in 4% paraformaldehyde
Negative Control:	One control slice for each experimental procedure processed omitting the primary antibody; overnight incubation in diluent solution only.
Notes:	The beta Amyloid antibody (Amyloid beta) (C-Term) ABIN5508784 works in IHC-P, especially at higher concentrations (1:200 and 1:500). Tyramide amplification strongly enhances the positive signal.
Primary Antibody:	ABIN5508784
Secondary Antibody:	poly-HRP conjugated goat anti-rabbit antibody
Protocol:	<ul style="list-style-type: none">• Perfuse mice with paraformaldehyde 4% in 0.1 M phosphate buffer pH 7.4 and post-fix in the same fixative for an additional 2 h at RT.• Wash, dehydrate, and embed samples in paraffin wax.• Wash several times with 0.01 M PBS.• Cut intestines and brain with a microtome into 6 µm sections and mount on glass slides.• After paraffin removal, incubate sections for 1 h at RT in PBS containing 1% albumin from chicken egg white (Sigma, A5378) and 0.3% Triton-X-100 (BioRad, 161-0407, lot 00583) to block non-specific binding sites.• Incubate sections with primary anti-beta amyloid (N-Term) (antibodies-online, ABIN5508784, lot 190522A) diluted 1:200, 1:500, 1:1000, and 1:2000 in 0.1 M PBS-BSA-PLL ON at RT.• Wash sections 3x 5 min with 0.01 M PBS.• Incubate sections with poly-HRP conjugated goat anti-rabbit antibody for 1 h at RT.• Wash sections 3x 5 min with 0.01 M PBS.• Incubate sections Tyramide working solution (for 5 sections: 100X Tyramide stock solution 5 µL, 100X H₂O₂ solution 5 µL, and 1X Reaction buffer 500 µL; Thermo Fisher Scientific, B40915, lot 2086865) for 10 min.• Stop the reaction with the Reaction Stop Reagent working solution.• Wash sections 3x 5 min with 0.01M PBS.

- Mount specimens in Fluoroshield (Sigma, F6182, lot MKCB0153V).
 - Acquire images with a fluorescence microscope and appropriate filter settings for AF488, e.g. Leica DM 6000B fluorescence microscope equipped with a digital camera at 40x magnification.
-

Image for Validation report #104493



Validation image no. 1 for anti-Amyloid beta (Abeta) (C-Term) antibody (ABIN5508784)

Staining of beta-amyloid positive neurons in the hippocampus and cerebral cortex of a 3-month-old transgenic mouse using ABIN5508784. Tyramide amplification. No counterstain.