antibodies -online.com







FGF10 Protein

Images



Overview

| Quantity: | 25 μg |
|----------------------|----------------------------|
| Target: | FGF10 |
| Origin: | Human |
| Source: | Escherichia coli (E. coli) |
| Protein Type: | Recombinant |
| Biological Activity: | Active |
| Application: | Functional Studies (Func) |

Product Details

| Characteristics: | Recombinant Human FGF10 |
|------------------|-------------------------------------|
| Purity: | >95 % as determined by SDS-PAGE |
| Endotoxin Level: | < 1.0 EU of endotoxin/µg of protein |

Target Details

Pathways:

| Target: | FGF10 |
|-------------------|------------------------|
| Alternative Name: | FGF10 (FGF10 Products) |
| Molecular Weight: | 19 |
| Gene ID: | 2255 |
| UniProt: | 015520 |

 $Order\ at\ www.antibodies-online.com\ |\ www.antiboerper-online.de\ |\ www.anticorps-enligne.fr\ |\ www.antibodies-online.com\ |\ www.antiboerper-online.de\ |\ www.antiboerper-online.d$ International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5509306 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

RTK Signaling, Fc-epsilon Receptor Signaling Pathway, EGFR Signaling Pathway, Neurotrophin


Signaling Pathway, Stem Cell Maintenance, Tube Formation, Positive Regulation of Response to DNA Damage Stimulus

Application Details


| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Restrictions: | For Research Use only |

| Handling | |
|------------------|---|
| Format: | Lyophilized |
| Reconstitution: | A quick spin of the vial followed by reconstitution in sterile distilled water to a concentration not less than 0.1 mg/mL is recommended. Please note, filter sterilization is a must following reconstitution. This solution can then be diluted into other buffers. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | The lyophilized protein is stable for at least one year from date of receipt at -70°C. Upon reconstitution, this cytokine can be stored in working aliquots at 2° - 8°C for one month, or at -20°C for six months, with a carrier protein without detectable loss of activity. Avoid repeated freeze/thaw cycles. |

Validation report #103788 for Haemagglutination (H)



SDS-PAGE Image 1.



Activity Assay

Image 2.