# .-online.com antibodies

# Datasheet for ABIN5510614 TGFB3 ELISA Kit

Image



#### Overview

Quantity:	96 tests
Target:	TGFB3
Binding Specificity:	AA 301-412
Reactivity:	Cow
Method Type:	Sandwich ELISA
Application:	ELISA

### Product Details

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of activated Bovine TGF-beta 3
Brand:	PicoKine™
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	Expression system for standard: sf21 Immunogen sequence: A301-S412
Cross-Reactivity (Details):	There is no detectable cross-reactivity with other relevant proteins.
Characteristics:	Sequence similarities: Belongs to the TGF-beta family.

## Target Details

Target:	TGFB3
Alternative Name:	TGFB3 (TGFB3 Products)


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5510614 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

Target Details	
Background:	Protein Function: Involved in embryogenesis and cell differentiation.
	Background: Transforming growth factor beta 3(TGF-beta 3) is a type of protein, known as a
	cytokine, which is involved in cell differentiation, embryogenesis and development. It belongs to
	a large family of cytokines called the Transforming growth factor beta superfamily. TGF-beta 3
	is believed to regulate molecules involved in cellular adhesion and extracellular matrix(ECM)
	formation during the process of palate development. Without TGF-beta 3, mammals develop a
	deformity known as acleft palate. This is caused by failure of epithelial cells in both sides of the
	developing palate to fuse. TGF-beta 3 also plays an essential role in controlling the
	development of lungs in mammals, by also regulating cell adhesion and ECM formation in this
	tissue, and controls wound healing by regulating the movements of epidermal and dermal cells
	in injured skin. TGF-beta 3 activated Lef1 in the absence of beta-catenin(CTNNB1) via nuclear
	phospho-Smad2 and Smad4.
	Synonyms: Transforming growth factor beta-3,TGF-beta-3,Latency-associated
	peptide,LAP,TGFB3,
	Full Gene Name: Transforming growth factor beta-3
	Cellular Localisation: Secreted.
UniProt:	P10600
Pathways:	Cell-Cell Junction Organization, Production of Molecular Mediator of Immune Response, Protein targeting to Nucleus

## Application Details

Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C,-20 °C
Storage Comment:	Store at 4 °C for 6 months, at -20 °C for 12 months. Avoid multiple freeze-thaw cycles(Shipped with wet ice.)
Expiry Date:	12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5510614 | 09/10/2023 | Copyright antibodies-online. All rights reserved.



#### ELISA

**Image 1.** Bovine TGF-beta 3 PicoKine ELISA Kit standard curve

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN5510614 | 09/10/2023 | Copyright antibodies-online. All rights reserved.