.-online.com antibodies

Datasheet for ABIN5510705

Image



Overview

Quantity:	96 tests
Target:	LDLR
Binding Specificity:	AA 22-790
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Sandwich High Sensitivity ELISA kit for Quantitative Detection of Mouse LDLR	
Brand:	PicoKine™	
Analytical Method:	Quantitative	
Detection Method:	Colorimetric	
Specificity:	Expression system for standard: NSO Immunogen sequence: A22-R790	
Cross-Reactivity (Details):	There is no detectable cross-reactivity with other relevant proteins.	

Target Details

Target:	LDLR
Alternative Name:	Ldlr (LDLR Products)
Background:	Protein Function: Binds LDL, the major cholesterol-carrying lipoprotein of plasma, and

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN5510705 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

UniProt:	P35951
	presence of PCSK9 (By similarity)
	Localizes to the Golgi apparatus, early and late endosomes/lysosomes and cell surface in the
	Lysosome . Found distributed from the plasma membrane to intracellular compartments.
	membrane protein . Golgi apparatus . Early endosome . Late endosome . Cell surface .
	system, Single-pass type I membrane protein. Membrane, clathrin-coated pit, Single-pass type I
	Cellular Localisation: Cell membrane, Single-pass type I membrane protein. Endomembrane
	Full Gene Name: Low-density lipoprotein receptor
	Synonyms: Low-density lipoprotein receptor,LDL receptor,Ldlr,
	Alternate splicing results in multiple transcript variants.
	Mutations in this gene cause the autosomal dominant disorder, familial hypercholesterolemia.
	synthesis. At the same time, a reciprocal stimulation of cholesterol ester synthesis takes place.
	methylglutaryl coenzyme A (HMG CoA) reductase, the rate-limiting step in cholesterol
	the cholesterol is made available for repression of microsomal enzyme 3-hydroxy-3-
	membrane and taken into the cell ending up in lysosomes where the protein is degraded and
	endocytosis of specific ligands. Low density lipoprotein (LDL) is normally bound at the cell
	receptor (LDLR) gene family consists of cell surface proteins involved in receptor-mediated
	belongs to the Low density lipoprotein receptor gene family. The low density lipoprotein
	cholesterol-rich LDL. In humans, the LDL receptor protein is encoded by the LDLR gene. It
	acids (after removal of 21-amino acid signal peptide) that mediates the endocytosis of
	Background: The Low-Density Lipoprotein (LDL) Receptor is a mosaic protein of 839 amino
	must first cluster into clathrin-coated pits.
	transports it into cens by endocytosis. In order to be internalized, the receptor light complexes

transports it into cells by endocytosis. In order to be internalized, the receptor-ligand complexes

Pathways:

Hepatitis C, Lipid Metabolism

Application Details

Plate:	Pre-coated	
Restrictions:	For Research Use only	
Handling		
Storage:	4 °C,-20 °C	
Storage Comment:	Store at 4°C for 6 months, at -20°C for 12 months. Avoid multiple freeze-thaw cycles(Shipped with wet ice.)	


Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN5510705 | 09/10/2023 | Copyright antibodies-online. All rights reserved.

 	•
ond	lina
and	
and	III I G

Expiry Date:

12 months

Images



ELISA

Image 1. Mouse LDLR PicoKine ELISA Kit standard curve