

## Datasheet for ABIN5516035

## anti-GNRHR antibody (C-Term)



Go to Product page

| $\sim$ |     |   |    |     |     |
|--------|-----|---|----|-----|-----|
|        | 1// | Р | rv | I P | ۱۸/ |
|        |     |   |    |     |     |

| Quantity:                      | 100 μL   |  |
|--------------------------------|--|--|
| Target:                        | GNRHR  |  |
| Binding Specificity:           | C-Term   |  |
| Reactivity:                    | Human  |  |
| Host:                          | Rabbit   |  |
| Clonality:                     | Polyclonal   |  |
| Conjugate:                     | This GNRHR antibody is un-conjugated   |  |
| Application:                   | Western Blotting (WB)  |  |
| Product Details                |  |  |
| Immunogen:                     | The immunogen is a synthetic peptide directed towards the C terminal region of human GNRHR   |  |
| Sequence:                      | ICNAKIIFTL TRVLHQDPHE LQLNQSKNNI PRARLKTLKM TVAFATSFTV   |  |
| Purification:                  | Affinity purified  |  |
| Target Details                 |  |  |
| Target:                        | GNRHR  |  |
|                                | GNRHR (GNRHR Products)   |  |
| Alternative Name:              | GNRHR (GNRHR Products)   |  |
| Alternative Name:  Background: | GNRHR (GNRHR Products)  This gene encodes the receptor for type 1 gonadotropin-releasing hormone. This receptor is a member of the seven-transmembrane, G-protein coupled receptor (GPCR) family. It is expressed on the surface of pituitary gonadotrope cells as well as lymphocytes, breast, ovary, |  |

and prostate. Following binding of gonadotropin-releasing hormone, the receptor associates with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Activation of the receptor ultimately causes the release of gonadotropic luteinizing hormone (LH) and follicle stimulating hormone (FSH). Defects in this gene are a cause of hypogonadotropic hypogonadism (HH). Alternative splicing results in multiple transcript variants encoding different isoforms. More than 18 transcription initiation sites in the 5' region and multiple polyA signals in the 3' region have been identified for this gene.

Alias Symbols: HH7, GRHR, LRHR, LHRHR, GNRHR1

Protein Size: 328

 Gene ID:
 2798

 NCBI Accession:
 NM\_000406, NP\_000397

 UniProt:
 P30968

Optimal working dilution should be determined by the investigator.

## **Application Details**

Application Notes:

| Restrictions:      | For Research Use only   |  |
|--------------------|---|--|
| Handling           |   |  |
| Format:            | Liquid  |  |
| Buffer:            | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.                                     |  |
| Preservative:      | Sodium azide  |  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.                  |  |
| Storage:           | -20 °C  |  |
| Storage Comment:   | For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. |  |