

# Datasheet for ABIN5517321

# anti-SETD7 antibody (N-Term)

# 1 Image



Go to Product page

| _ |     |          |    |    |    |    |
|---|-----|----------|----|----|----|----|
|   | 1// | Д        | rv | 16 | ٦/ | ٨  |
| U | W   | $\vdash$ | ΙV | Ιt | ٦, | /V |

| Quantity:             | 100 μL   |
|-----------------------|--|
| Target:               | SETD7  |
| Binding Specificity:  | N-Term   |
| Reactivity:           | Human, Mouse, Rat, Dog, Horse, Pig, Rabbit, Cow, Zebrafish (Danio rerio), Guinea Pig   |
| Host:                 | Rabbit   |
| Clonality:            | Polyclonal   |
| Conjugate:            | This SETD7 antibody is un-conjugated   |
| Application:          | Western Blotting (WB)  |
| Product Details       |  |
| Immunogen:            | The immunogen is a synthetic peptide directed towards the N-terminal region of Human SETD7                                     |
| Sequence:             | MDSDDEMVEE AVEGHLDDDG LPHGFCTVTY SSTDRFEGNF VHGEKNGRGK   |
| Predicted Reactivity: | Cow: 93%, Dog: 93%, Guinea Pig: 86%, Horse: 100%, Human: 100%, Mouse: 100%, Pig: 100%, Rabbit: 100%, Rat: 100%, Zebrafish: 92% |
| Characteristics:      | This is a rabbit polyclonal antibody against SETD7. It was validated on Western Blot.  |
| Purification:         | Affinity purified  |
| Target Details        |  |
| Target:               | SETD7  |
|                       |  |
| Alternative Name:     | SETD7 (SETD7 Products)   |

#### **Target Details**

Background:

SETD7 is a histone methyltransferase that specifically monomethylates 'Lys-4' of histone H3. H3 'Lys-4' methylation represents a specific tag for epigenetic transcriptional activation. It plays a central role in the transcriptional activation of genes such as collagenase or insulin. Recruited by IPF1/PDX-1 to the insulin promoter, leading to activate transcription. Has also methyltransferase activity toward non-histone proteins such as p53/TP53, TAF10, and possibly TAF7 by recognizing and binding the [KR]-[STA]-K in substrate proteins. Monomethylates 'Lys-189' of TAF10, leading to increase the affinity of TAF10 for RNA polymerase II. Monomethylates 'Lys-372' of p53/TP53, stabilizing p53/TP53 and increasing p53/TP53-mediated transcriptional activation.

Alias Symbols: SETD7, KIAA1717, KMT7, SET7, SET9,

Protein Interaction Partner: KLF4, HIST3H3, SOX2, PRKAR1A, A1BG, TP53, NR2F2, STEAP3, SERTAD1, DPY30, MED1, MYOD1, E2F1, HIST1H3A, SMARCA4, H3F3A, STAT3, RELA, AR, UBC, DNMT1, PPP1R12A, SIRT1, TBP, TAF10, TAF7, TAF6, HIST1H4A, ESR1, HIST2H3C, HCFC1, L3MBTL1,

Protein Size: 366

Q8WTS6

 Gene ID:
 80854

 NCBI Accession:
 NP\_085151

#### **Application Details**

**Application Notes:** 

UniProt:


| Restrictions:      | For Research Use only   |
|--------------------|---|
| Handling           |   |
| Format:            | Liquid  |
| Buffer:            | Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose. |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOLIS AND HAZARDOLIS SUBSTANCE which                     |

Optimal working dilution should be determined by the investigator.

## Handling

|                  | should be handled by trained staff only.  |
|------------------|---|
| Storage:         | -20 °C  |
| Storage Comment: | For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles. |

### **Images**



### **Western Blotting**

Image 1. Host: Rabbit Target Name: SETD7 Sample Type: Jurkat Whole Cell lysates Antibody Dilution: 1.0ug/ml