

[Go to Product page](#)

## Datasheet for ABIN5518102 **anti-CIRH1A antibody (Middle Region)**

### Overview

Quantity:	100 µL
Target:	CIRH1A
Binding Specificity:	Middle Region
Reactivity:	Human, Mouse, Rat, Horse, Rabbit, Cow, Dog, Guinea Pig
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB)

### Product Details

Immunogen:	The immunogen for Anti-CIRH1A antibody is: synthetic peptide directed towards the Middle region of Human CIR1A
Sequence:	FAHHLELWRL GSTVATGKNG DTLPLSKNAD HLLHLKTKGP ENICSCISP
Predicted Reactivity:	Cow: 100%, Dog: 100%, Guinea Pig: 100%, Horse: 100%, Human: 100%, Mouse: 100%, Rabbit: 100%, Rat: 100%
Characteristics:	This is a rabbit polyclonal antibody against CIR1A. It was validated on Western Blot.
Purification:	Affinity purified

### Target Details

Target:	CIRH1A
Alternative Name:	CIRH1A ( <a href="#">CIRH1A Products</a> )

## Target Details

Background:	<p>This gene encodes a WD40-repeat-containing protein that is localized to the nucleolus.</p> <p>Mutation of this gene causes North American Indian childhood cirrhosis, a severe intrahepatic cholestasis that results in transient neonatal jaundice, and progresses to periportal fibrosis and cirrhosis in childhood and adolescence.</p> <p>Alias Symbols: NAIC, UTP4, CIRHIN, TEX292</p> <p>Protein Interaction Partner: UBC, SUMO1, NEDD8, RNF2, HDAC11, UTP15, GLTSCR2, APP, UBD, PRNP, ISG15, SIRT7,</p> <p>Protein Size: 625</p>
-------------	--

Gene ID:	84916
----------	-------

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Buffer:	Liquid. Purified antibody supplied in 1x PBS buffer with 0.09 % (w/v) sodium azide and 2 % sucrose.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	For short term use, store at 2-8°C up to 1 week. For long term storage, store at -20°C in small aliquots to prevent freeze-thaw cycles.