# antibodies .- online.com





anti-GAS6 antibody (AA 488-660)





Overview	
Quantity:	100 μg
Target:	GAS6
Binding Specificity:	AA 488-660
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GAS6 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Rabbit IgG polyclonal antibody for Growth arrest-specific protein 6(GAS6) detection. Tested with WB, IHC-P in Human, Mouse, Rat.
lmmunogen:	E.coli-derived human GAS 6 recombinant protein (Position: M488-S660). Human GAS 6 shares 79.7% and 81.4% amino acid (aa) sequence identity with mouse and rat GAS 6, respectively.
Isotype:	IgG
Cross-Reactivity (Details):	Predicted Cross Reactivity: human  No cross reactivity with other proteins.  Predicted Cross Reactivity: Species predicted to be fit for the product based on sequence similarities.
Characteristics:	Rabbit IgG polyclonal antibody for Growth arrest-specific protein 6(GAS6) detection. Tested with WB, IHC-P in Human, Mouse, Rat.


# **Product Details** Gene Name: growth arrest specific 6 Protein Name: Growth arrest-specific protein 6 Purification: Immunogen affinity purified. **Target Details** GAS<sub>6</sub> Target: Alternative Name: GAS6 (GAS6 Products) Background: Growth arrest-specific 6, also known as GAS6, is a human gene coding for the Gas6 protein. It is similar to the Protein S with the same domain organization and 43 % amino acid identity. It was originally found as a gene upregulated by growth arrested fibroblasts. This gene is frequently overexpressed in many cancers and has been implicated as an adverse prognostic marker. Elevated protein levels are additionally associated with a variety of disease states, including venous thromboembolic disease, systemic lupus erythematosus, chronic renal failure, and preeclampsia. Synonyms: AXLLG | AXS F | AXSF | FLJ34709 | Gas 6 | GAS-6 | Gas6 | Q14393 Gene ID: 2621 UniProt: Q14393 Pathways: RTK Signaling, Carbohydrate Homeostasis, Production of Molecular Mediator of Immune Response **Application Details**

Application Notes:	WB: Concentration: 0.1-0.5 μg/mL, Tested Species: Mouse, Rat, Predicted Species: Human
	IHC-P: Concentration: 0.5-1 μg/mL, Tested Species: Mouse, Rat, Predicted Species: Human,
	Epitope Retrieval by Heat: Boiling the paraffin sections in 10 mM citrate buffer, pH 6.0, for
	20 mins is required for the staining of formalin/paraffin sections.
	Notes: Tested Species: Species with positive results. Predicted Species: Species predicted to be
	fit for the product based on sequence similarities. Other applications have not been tested.
	Optimal dilutions should be determined by end users.
Comment:	Antibody can be supported by chemiluminescence kit ABIN921124 in WB, supported by
	ABIN921231 in IHC(P).
Restrictions:	For Research Use only

## Handling


Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 $\mu g/mL$ .
Concentration:	500 μg/mL
Buffer:	Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month.  It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

# **Images**



#### **Western Blotting**


Image 1. Western blot analysis of GAS 6 using anti-GAS 6 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat brain tissue lysates, Lane 2: mouse brain tissue lysates. After Electrophoresis, proteins were transferred to Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GAS 6 antigen affinity purified polyclonal antibody (Catalog # ) at 0.5  $\mu$ g/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for GAS 6 at approximately 75KD. The expected



band size for GAS 6 is at 79KD.

#### **Immunohistochemistry**

**Image 2.** IHC analysis of GAS 6 using anti- GAS 6 antibody . GAS 6 was detected in paraffin-embedded section of mouse spleen tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml rabbit anti- GAS 6 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



### **Immunohistochemistry**

**Image 3.** IHC analysis of GAS 6 using anti- GAS 6 antibody . GAS 6 was detected in paraffin-embedded section of rat spleen tissues. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1 $\mu$ g/ml rabbit anti- GAS 6 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.