# antibodies -online.com







# anti-DLG3 antibody (AA 749-817)

**Images** 



| Coto  | Product | nage |
|-------|---------|------|
| GU 10 | FIUUUUU | paye |

| Overview                    |  |  |
|-----------------------------|--|--|
| Quantity:                   | 100 μg   |  |
| Target:                     | DLG3   |  |
| Binding Specificity:        | AA 749-817   |  |
| Reactivity:                 | Rat, Mouse   |  |
| Host:                       | Rabbit   |  |
| Clonality:                  | Polyclonal   |  |
| Conjugate:                  | This DLG3 antibody is un-conjugated  |  |
| Application:                | Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))   |  |
| Product Details             |  |  |
| Purpose:                    | Rabbit IgG polyclonal antibody for Disks large homolog 3(DLG3) detection. Tested with WB, IHC-P in Human,Mouse,Rat.  |  |
| Immunogen:                  | E. coli-derived human SAP102 recombinant protein (Position: E749-L817). Human SAP102 shares 98.6% amino acid (aa) sequence identity with both mouse and rat SAP102.                  |  |
| Isotype:                    | IgG  |  |
| Cross-Reactivity (Details): | Predicted Cross Reactivity: human  No cross reactivity with other proteins.  Predicted Cross Reactivity: Species predicted to be fit for the product based on sequence similarities. |  |
| Characteristics:            | Rabbit IgG polyclonal antibody for Disks large homolog 3(DLG3) detection. Tested with WB, IHC-P in Human,Mouse,Rat.  |  |

| Product Details     |   |  |
|---------------------|---|--|
|                     | Gene Name: discs large MAGUK scaffold protein 3   |  |
|                     | Protein Name: Disks large homolog 3   |  |
| Purification:       | Immunogen affinity purified.  |  |
| Target Details      |   |  |
| Target:             | DLG3  |  |
| Alternative Name:   | DLG3 (DLG3 Products)  |  |
| Background:         | Disks large homolog 3 (DLG3) also known as neuroendocrine-DLG or synapse-associated protein 102 (SAP-102) is a protein that in humans is encoded by the DLG3 gene. DLG3 is a member of the membrane-associated guanylate kinase (MAGUK) superfamily of proteins. The encoded protein may play a role in clustering of NMDA receptors at excitatory synapses. It may also negatively regulate cell proliferation through interaction with the C-terminal region of the adenomatosis polyposis coli tumor suppressor protein. Mutations in this gene have been associated with X-linked mental retardation. |  |
|                     | Synonyms: Disks large homolog 3, Neuroendocrine-DLG, Synapse-associated protein 102, SAP-102, SAP102, XLMR, DLG3, KIAA1232  |  |
| Gene ID:            | 1741  |  |
| UniProt:            | Q92796  |  |
| Application Details |   |  |
| Application Notes:  | WB: Concentration: 0.1-0.5 µg/mL, Tested Species: Mouse, Rat, Predicted Species: Human IHC-P: Concentration: 0.5-1 µg/mL, Tested Species: Mouse, Rat, Predicted Species: Human, Epitope Retrieval by Heat: Boiling the paraffin sections in 10 mM citrate buffer, pH 6.0, for 20 mins is required for the staining of formalin/paraffin sections.  Notes: Tested Species: Species with positive results. Predicted Species: Species predicted to be fit for the product based on sequence similarities. Other applications have not been tested. Optimal dilutions should be determined by end users.     |  |
| Comment:            | Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (ABIN921124) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for  |  |

IHC(P).


For Research Use only

Restrictions:

## Handling


| Format:            | Lyophilized   |
|--------------------|---|
| Reconstitution:    | Add 0.2 mL of distilled water will yield a concentration of 500 µg/mL.  |
| Concentration:     | 500 μg/mL   |
| Buffer:            | Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.   |
| Preservative:      | Sodium azide  |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.  |
| Storage:           | 4 °C,-20 °C   |
| Storage Comment:   | At -20°C for one year. After reconstitution, at 4°C for one month.  It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing. |


# **Images**



#### **Immunohistochemistry**

Image 1. IHC analysis of SAP102 using anti-SAP102 antibody . SAP102 was detected in paraffin-embedded section of mouse brain tissue . Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1μg/ml rabbit anti-SAP102 Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.





### **Immunohistochemistry**

Image 2. IHC analysis of SAP102 using anti-SAP102 antibody . SAP102 was detected in paraffin-embedded section of rat brain tissue . Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1µg/ml rabbit anti-SAP102 Antibody overnight at 4°C. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

#### **Western Blotting**

Image 3. Western blot analysis of SAP102 using anti-SAP102 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat brain tissue lysate, Lane 2: mouse brain tissue lysate. After Electrophoresis, transferred proteins were Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-SAP102 antigen affinity purified polyclonal antibody (Catalog # ) at 0.5 µg/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for SAP102 at approximately 102KD. The expected band size for SAP102 is at 90KD.