

# Datasheet for ABIN5518996

# anti-GSTP1 antibody (AA 2-210)



Overview

Purification:

3

**Publications** 



Go to Product page

Quantity:	100 μg
Target:	GSTP1
Binding Specificity:	AA 2-210
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GSTP1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Rabbit IgG polyclonal antibody for GST3 / GST pi detection. Tested with WB, IHC-P, Direct ELISA in Human,Mouse,Rat.
Immunogen:	E. coli-derived mouse GST3 / GST pi recombinant protein (Position: P2-Q210).
Isotype:	IgG
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Characteristics:	Rabbit IgG polyclonal antibody for GST3 / GST pi detection. Tested with WB, IHC-P, Direct ELISA in Human, Mouse, Rat.  Gene Name: glutathione S-transferase, pi 1

Protein Name: Glutathione S-transferase P 1

Immunogen affinity purified.

# **Target Details**

Target:	GSTP1
Alternative Name:	Gstp1 (GSTP1 Products)
Background:	Glutathione S-transferases pi (GSTP1), also known as GST3, is an enzyme that in humans is encoded by the GSTP1 gene. This gene is mapped to 11q13.2. GSTP1 has 7 exons and 6 introns contained within approximately 2.8 kilobases. GSTP1 belongs to Glutathione S-transferases (GSTs) which are a family of enzymes that play an important role in detoxification by catalyzing the conjugation of many hydrophobic and electrophilic compounds with reduced glutathione. GSTP1 presents in all tissues and cells, with the exception of red cells, in which only erythrocyte GST(GSTe) is observed. What's more, GSTP1 is a polymorphic gene encoding active, functionally different GSTP1 variant proteins that are thought to function in xenobiotic metabolism and play a role in susceptibility to cancer, and other diseases.
	Synonyms: Glutathione S-transferase P 1, Gst P1, GST YF-YF, GST class-pi, GST-piB, Preadipocyte growth factor, Gstp1, Gstpib
Gene ID:	14870
UniProt:	P19157
Pathways:	Cellular Response to Molecule of Bacterial Origin
Application Details	
Application Notes:	Notes: Tested Species: Species with positive results. Other applications have not been tested.  Optimal dilutions should be determined by end users.
Comment:	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (ABIN921124) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 μg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.

# Handling

Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month.  It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

#### **Publications**


#### Product cited in:

Zhang, Yan, Xu, Fang, Zhang, Zhang, Wu, Kong, Huang: "The recovery trajectory of adolescent social defeat stress-induced behavioral, (1)H-MRS metabolites and myelin changes in Balb/c mice." in: **Scientific reports**, Vol. 6, pp. 27906, (2018) (PubMed).

Sun, Hu, Xiong, Tu, Zhao, Chen, Song, Ye: "Enhanced expression of stem cell markers and drug resistance in sphere-forming non-small cell lung cancer cells." in: **International journal of clinical and experimental pathology**, Vol. 8, Issue 6, pp. 6287-300, (2016) (PubMed).

Yang, Gao, Zhang, Fang, Wu, Xu, Huang: "Changes in proinflammatory cytokines and white matter in chronically stressed rats." in: **Neuropsychiatric disease and treatment**, Vol. 11, pp. 597-607, (2015) (PubMed).


#### **Images**



## **Western Blotting**


Image 1. Western blot analysis of GST3 / GST pi using anti-GST3 / GST pi antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: mouse lung tissue lysates, Lane 2: mouse testis tissue lysates, Lane 3: rat spleen tissue lysates, Lane 4: human A549 whole cell lysates, Lane 5: human placenta tissue lysates, Lane 6: human Hela whole cell lysates. After

Electrophoresis, proteins were transferred to Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-GST3 / GST pi antigen affinity purified polyclonal antibody (Catalog #) at 0.5 ug/mL overnight at 4°C, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for GST3 / GST pi at approximately 23KD. The expected band size for GST3 / GST pi is at 23KD.



#### **Immunohistochemistry**

Image 2. IHC analysis of GST3 / GST pi using anti-GST3 / GST pi antibody .GST3 / GST pi was detected in paraffinembedded section of mouse kidney tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-GST3 / GST pi Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



## **Immunohistochemistry**

Image 3. IHC analysis of GST3 / GST pi using anti-GST3 / GST pi antibody .GST3 / GST pi was detected in paraffinembedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-GST3 / GST pi Antibody overnight at 4°C. Biotinylated goat anti-rabbit lgG was used as secondary antibody and incubated for 30 minutes at 37°C. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Please check the product details page for more images. Overall 5 images are available for ABIN5518996.