antibodies -online.com







anti-PARK7/DJ1 antibody (AA 2-189)



Images



Go to	Prod	uct	page

()	VE	۲۷	ΊE	W
\circ	V	- I V	10	, v v

Overview	
Quantity:	100 μg
Target:	PARK7/DJ1 (PARK7)
Binding Specificity:	AA 2-189
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Purpose:	Rabbit IgG polyclonal antibody for PARK7 / DJ1 detection. Tested with WB, IHC-P, Direct ELISA in Mouse,Rat.
Purpose: Immunogen:	
	in Mouse,Rat.
Immunogen:	in Mouse,Rat. E. coli-derived rat PARK7 / DJ1 recombinant protein (Position: A2-D189).
Immunogen: Isotype:	in Mouse,Rat. E. coli-derived rat PARK7 / DJ1 recombinant protein (Position: A2-D189). IgG


Target Details

Target:	PARK7/DJ1 (PARK7)
Alternative Name:	Park7 (PARK7 Products)
Background:	Parkinson disease (autosomal recessive, early onset) 7, also known as DJ1, is a protein which in humans is encoded by the PARK7 gene. PARK7 belongs to the peptidase C56 family of proteins. PARK7 is mapped to chromosome 1p36. It acts as a positive regulator of androgen receptor-dependent transcription. It is also involved in tumorigenesis and in maintaining mitochondrial homeostasis. This gene may also function as a redox-sensitive chaperone, as a sensor foroxidative stress, and it apparently protects neurons against oxidative stress and cell death. It has been found that PARK7 mutations that impair transcriptional coactivator function can render dopaminergic neurons vulnerable to apoptosis and may contribute to the pathogenesis of Parkinson disease.
Gene ID:	Synonyms: Protein DJ-1, DJ-1, Contraception-associated protein 1 117287
UniProt:	088767
Pathways:	Intracellular Steroid Hormone Receptor Signaling Pathway, Regulation of Intracellular Steroid Hormone Receptor Signaling, Proton Transport
Application Details	
Application Notes:	Notes: Tested Species: Species with positive results. Other applications have not been tested. Optimal dilutions should be determined by end users.
Comment:	Boster recommends Enhanced Chemiluminescent Kit with anti-Rabbit IgG (ABIN921124) for Western blot, and HRP Conjugated anti-Rabbit IgG Super Vision Assay Kit (SV0002-1) for IHC(P).
Restrictions:	For Research Use only
Handling	
Format:	Lyophilized
Reconstitution:	Add 0.2 mL of distilled water will yield a concentration of 500 μg/mL.
Concentration:	500 μg/mL
Buffer:	Each vial contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2HPO4, 0.05 mg Sodium azide.

Handling


Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for one year. After reconstitution, at 4°C for one month. It can also be aliquotted and stored frozen at -20 °C for a longer time. Avoid repeated freezing and thawing.

Images



Immunohistochemistry


Image 1. IHC analysis of PARK7 / DJ1 using anti-PARK7 / DJ1 antibody .PARK7 / DJ1 was detected in paraffinembedded section of rat testis tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-PARK7 / DJ1 Antibody overnight at 4â, f. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37â, f. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.



Western Blotting

Image 2. Western blot analysis of PARK7 / DJ1 using anti-PARK7 / DJ1 antibody. Electrophoresis was performed on a 5-20% SDS-PAGE gel at 70V (Stacking gel) / 90V (Resolving gel) for 2-3 hours. The sample well of each Lane was loaded with 50ug of sample under reducing conditions. Lane 1: rat pancreas tissue lysates, Lane 2: rat kidney tissue lysates, Lane 3: rat skeletal muscle tissue lysates, Lane 4: rat liver tissue lysates, Lane 5: rat testis tissue lysates, Lane 6: rat heart tissue lysates, Lane 7: mouse kidney tissue lysates,

Lane 8: mouse skeletal muscle tissue lysates. After Electrophoresis, proteins were transferred to Nitrocellulose membrane at 150mA for 50-90 minutes. Blocked the membrane with 5% Non-fat Milk/ TBS for 1.5 hour at RT. The membrane was incubated with rabbit anti-PARK7 / DJ1 antigen affinity purified polyclonal antibody (Catalog #) at 0.5 ug/mL overnight at 4â, f, then washed with TBS-0.1%Tween 3 times with 5 minutes each and probed with a goat anti-rabbit IgG-HRP secondary antibody at a dilution of 1:10000 for 1.5 hour at RT. The signal is developed using an Enhanced Chemiluminescent detection (ECL) kit (Catalog # EK1002) with Tanon 5200 system. A specific band was detected for PARK7 / DJ1 at approximately 22KD. The expected band size for PARK7 / DJ1 is at 20KD.



Immunohistochemistry

Image 3. IHC analysis of PARK7 / DJ1 using anti-PARK7 / DJ1 antibody .PARK7 / DJ1 was detected in paraffinembedded section of mouse liver tissue. Heat mediated antigen retrieval was performed in citrate buffer (pH6, epitope retrieval solution) for 20 mins. The tissue section was blocked with 10% goat serum. The tissue section was then incubated with 1ug/ml rabbit anti-PARK7 / DJ1 Antibody overnight at 4â, f. Biotinylated goat anti-rabbit IgG was used as secondary antibody and incubated for 30 minutes at 37â, f. The tissue section was developed using Strepavidin-Biotin-Complex (SABC)(Catalog # SA1022) with DAB as the chromogen.

Please check the product details page for more images. Overall 7 images are available for ABIN5519050.