

Datasheet for ABIN5521961

ENPP2 ELISA Kit





Overview

| Quantity: | 96 tests |
|--------------------------|-----------------------|
| Target: | ENPP2 |
| Reactivity: | Human |
| Method Type: | Sandwich ELISA |
| Detection Range: | 1.25 ng/mL - 80 ng/mL |
| Minimum Detection Limit: | 1.25 ng/mL |
| Application: | ELISA |

Product Details

| Troduct Details | |
|--------------------|--|
| Purpose: | For quantitative detection of ENPP2 in serum, plasma, tissue homogenates. |
| Sample Type: | Plasma, Serum, Tissue Homogenate |
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Specificity: | This assay has high sensitivity and excellent specificity for detection of ENPP2. No significant cross-reactivity or interference between ENPP2 and analogues was observed. Note: Limited by current skills and knowledge, it is difficult for us to complete the cross-reactivity detection between ENPP2 and all the analogues, therefore, cross reaction may still exist. |
| Sensitivity: | 0.75 ng/mL |
| Components: | - ELISA Microplate - Lyophilized Standard - Sample / Standard Dilution Buffer |

- Biotin-labeled Antibody (Concentrated)
- Antibody Dilution Buffer
- HRP-Streptavidin Conjugate(SABC)
- SABC Dilution Buffer
- TMB Substrate
- Stop Solution
- Wash Buffer (25X)
- Plate Sealer
- Product Description

Target Details

| Target: | ENPP2 |
|-------------------|---|
| Alternative Name: | ENPP2 |
| Background: | ENPP2/E-NPP 2/Autotaxin/ATX/Autotaxin/Lysophosphatidic Acid/NPP2/PDNP2/ATX-X/AUTOTAXIN/autotaxin-t/ectonucleotide pyrophosphatase/phosphodiesterase 2/Extracellular lysophospholipase D/LysoPLD/PD-IALPHA/phosphodiesterase I/nucleotide pyrophosphatase 2/plas |
| UniProt: | Q13822 |

Application Details

| Application Notes: | Optimal working dilution should be determined by the investigator. |
|--------------------|--|
| Comment: | Sandwich ELISA, Double Antibody |
| Sample Volume: | 100 μL |
| Plate: | Pre-coated |
| Restrictions: | For Research Use only |


| according to the labels on vials. The hould be stored at -20 °C upon receipt while eagents need to be stored according to the |
|---|
| re |

above storage condition. Besides, please return the unused wells to the foil pouch containing the desiccant pack, and zip-seal the foil pouch.

Expiry Date:

6 months

Images



ELISA

Image 1. A typical standard curve