

Datasheet for ABIN5526680
ULK1 ELISA Kit

3 Images

[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	ULK1
Binding Specificity:	pSer556
Reactivity:	Human, Mouse
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human and Mouse Phospho-ULK1 (S556) ELISA Kit. This assay semi-quantitatively measures ULK1 phosphorylated at Serine-556 in cell lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human and Mouse ULK1 phosphorylated at site Serine-556.
Characteristics:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer• Anti-Phospho Antibody• HRP-Conjugated Secondary Antibody• Assay Diluent• TMB One-Step Substrate• Stop Solution• Lysis Buffer

Product Details

- Positive Control Sample

Components:

- Pre-Coated 96-well Strip Microplate
- Wash Buffer
- Anti-Phospho Antibody
- HRP-Conjugated Secondary Antibody
- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target: ULK1

Alternative Name: ULK1 ([ULK1 Products](#))

Gene ID: 8408

UniProt: [O75385](#), [O70405](#)

Pathways: [Regulation of Cell Size](#), [Autophagy](#)

Application Details

Application Notes: Optimal working dilution should be determined by the investigator.

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.

Application Details

- 6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
- 7. Incubate 1 h at RT.
- 8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
- 9. Incubate 30 min at RT.
- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months

Images




ELISA

Image 1. Jurkat cells were treated with H2O2 at 37°C for 2 min. Cells were solubilized at 4 x 10⁷ cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.

ELISA

Image 2. C2C12 cells were untreated or treated with 1mM H2O2 for 15 min. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 3. C2C12 cells were untreated or treated with 1mM H2O2 for 15 min. Cell lysates were analyzed using this phosphoELISA and Western Blot.

