

Datasheet for ABIN5526693

**GATA2 ELISA Kit****2** Images[Go to Product page](#)

## Overview

Quantity:	96 tests
Target:	GATA2
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA

## Product Details

Purpose:	Human GATA-2 Transcription Factor Activity Assay. This assay uses a dsDNA coated plate with canonical GATA-2 binding sequences to semi-quantitatively detect active GATA-2 in lysates or nuclear extracts. Only available in North America.
Sample Type:	Cell Lysate, Nuclear Extract
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The oligonucleotide/antibody pair provided in this kit recognizes human FRA-2 in whole lysates and nuclear extracts.
Characteristics:	<ul style="list-style-type: none"><li>• Specific transcription factor-DNA binding assay</li><li>• Perfect alternative to EMSA</li><li>• Easy to perform in an ELISA format</li><li>• Non-radioactive assay</li><li>• High throughput (96 well plate format)</li><li>• Assay can be completed within 5 hours</li></ul>

## Product Details

Components:	<ul style="list-style-type: none"><li>• 96-well Strip Microplate pre-coated with DNA probes</li><li>• DNA Binding Buffer</li><li>• Positive Control Sample</li><li>• Specific Competitor DNA probe</li><li>• Non-specific Competitor DNA probe</li><li>• Assay Reagent</li><li>• DTT</li><li>• Wash Buffer</li><li>• Primary Antibody</li><li>• HRP-conjugated Secondary Antibody</li><li>• TMB One-Step Substrate Reagent</li><li>• Stop Solution</li></ul>
-------------	--

Material not included:	<ul style="list-style-type: none"><li>• Distilled or deionized water</li><li>• 100 mL and 1 liter graduated cylinders</li><li>• Tubes to prepare sample dilutions Absorbent paper</li><li>• Precision pipettes to deliver 2 µL to 1 mL volumes</li><li>• Adjustable 1-25 mL pipettes for reagent preparation</li><li>• Benchtop rocker or shaker</li><li>• Microplate reader capable of measuring absorbance at 450 nm</li></ul>
------------------------	--

## Target Details

Target:	GATA2
Alternative Name:	GATA-2 ( <a href="#">GATA2 Products</a> )
Gene ID:	2624
UniProt:	<a href="#">P23769</a>
Pathways:	<a href="#">Stem Cell Maintenance</a>

## Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none"><li>1. Prepare all reagents and samples as instructed in the manual.</li><li>2. Add 100 µL of sample or positive control to each well.</li><li>3. Incubate 2 h at RT or O/N at 4 °C.</li><li>4. Add 100 µL of prepared primary antibody to each well.</li><li>5. Incubate 1 h at RT.</li><li>6. Add 100 µL of prepared HRP-secondary antibody to each well.</li></ol>

Application Details

7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 °C

Storage Comment: Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw cycles.


Expiry Date: 6 months

Images



Activity Assay

**Image 1.** Transcription factor activity assay of GATA-2 from nuclear extracts of K562 cells or HeLa cells with the specific competitor or non-specific competitor. The result shows specific binding of GATA-2 to the GATA conserved binding site detected by using the GATA-2 TF-Activity Assay Kit.



Activity Assay

**Image 2.** Transcription factor activity assay of GATA-2 from nuclear extracts of K562 cells or HeLa cells. A. Western-blot result of GATA-2 from cytoplasmic and nuclear fractions. B. Transcription factor activity assay of GATA-2 from nuclear fractions with the GATA-2 TF-Activity Assay Kit.