

Datasheet for ABIN5526695

HDAC2 ELISA Kit**3** Images[Go to Product page](#)

Overview

Quantity:	96 tests
Target:	HDAC2
Binding Specificity:	pSer394
Reactivity:	Human, Mouse, Rat
Method Type:	Sandwich ELISA
Application:	ELISA

Product Details

Purpose:	Human, Mouse and Rat Phospho-HDAC2 (Ser394) ELISA Kit. This assay semi-quantitatively measures HDAC2 phosphorylated at Serine-394 in cell lysate samples.
Sample Type:	Cell Culture Lysate
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human, Mouse and Rat HDAC2 phosphorylated at site Serine-394.
Characteristics:	<ul style="list-style-type: none">• Rapidly measure phosphorylated protein in lysates• Screen numerous different cell lysates without performing a Western Blot analysis• Minimal hands-on time, convenient, and non-radioactive material
Components:	<ul style="list-style-type: none">• Pre-Coated 96-well Strip Microplate• Wash Buffer• Anti-Phospho Antibody• HRP-Conjugated Secondary Antibody

Product Details

- Assay Diluent
- TMB One-Step Substrate
- Stop Solution
- Lysis Buffer
- Positive Control Sample

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	HDAC2
Alternative Name:	HDAC2 (HDAC2 Products)
Gene ID:	3066
UniProt:	Q92769 , P70288 , F7ENH8
Pathways:	Neurotrophin Signaling Pathway , Regulation of Muscle Cell Differentiation , Negative Regulation of intrinsic apoptotic Signaling , SARS-CoV-2 Protein Interactome , The Global Phosphorylation Landscape of SARS-CoV-2 Infection

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none">1. Prepare all reagents and samples as instructed in the manual.2. Add 100 µL of sample or positive control to each well.3. Incubate 2.5 h at RT or O/N at 4 °C.4. Add 100 µL of prepared primary antibody to each well.5. Incubate 1 h at RT.6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.7. Incubate 1 h at RT.8. Add 100 µL of TMB One-Step Substrate Reagent to each well.9. Incubate 30 min at RT.

Application Details

- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

Restrictions: For Research Use only


Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months

Images




ELISA

Image 1. HeLa cells were treated with Calyculin A. Solubilize cells at 4×10^7 cells/ml in Cell Lysate Buffer. Serial dilutions of lysates were analyzed in this ELISA.

ELISA

Image 2. HeLa cells were treated or untreated with Calyculin A. A. Cell lysates were analyzed using this phosphoELISA and Western Blot.



ELISA

Image 3. HeLa cells were treated or untreated with Calyculin A. A. Cell lysates were analyzed using this phosphoELISA and Western Blot.

