antibodies .- online.com





Datasheet for ABIN5526700

NFYA ELISA Kit



Overview

Quantity:	96 tests
Target:	NFYA
Reactivity:	Human
Method Type:	DNA-Binding ELISA
Application:	ELISA
Product Details	
Purpose:	Human NF-YA Transcription Factor Activity Assay. This assay uses a dsDNA coated plate with canonical NF-YA binding sequences to semi-quantitatively detect active NF-YA in lysates or nuclear extracts. Only available in North America.
Sample Type:	Cell Lysate, Nuclear Extract
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	The olionucleotide/antibody pair provided in this kit recognizes mouse NFYA in whole lysates and nuclear extracts.
Characteristics:	 Specific transcription factor-DNA binding assay Perfect alternative to EMSA Easy to perform in an ELISA format Non-radioactive assay High throughput (96 well plate format) Assay can be completed within 5 hours

Product Details

Components:

- 96-well Strip Microplate pre-coated with DNA probes
- · DNA Binding Buffer
- · Positive Control Sample
- Specific Competitor DNA probe
- · Non-specific Competitor DNA probe
- · Assay Reagent
- DTT
- · Wash Buffer
- · Primary Antibody
- · HRP-conjugated Secondary Antibody
- TMB One-Step Substrate Reagent
- Stop Solution

Material not included:

- Distilled or deionized water
- · 100 mL and 1 liter graduated cylinders
- · Tubes to prepare sample dilutions Absorbent paper
- Precision pipettes to deliver 2 µL to 1 mL volumes
- · Adjustable 1-25 mL pipettes for reagent preparation
- · Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

Target Details

Target:	NFYA
Alternative Name:	NF-YA (NFYA Products)
Gene ID:	4800
UniProt:	P23511
Pathways:	Regulation of Lipid Metabolism by PPARalpha

Application Details	
Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	1. Prepare all reagents and samples as instructed in the manual.
	2. Add 100 μL of sample or positive control to each well.
	3. Incubate 2 h at RT or O/N at 4 °C.
	4. Add 100 μL of prepared primary antibody to each well.
	5. Incubate 1 h at RT.
	6. Add 100 μL of prepared HRP-secondary antibody to each well.

Application Details

	7. Incubate 1 h at RT. 8. Add 100 μL of TMB One-Step Substrate Reagent to each well.
	9. Incubate 30 min at RT.
	10. Add 50 μL of Stop Solution to each well.
	11. Read at 450 nm immediately.
Restrictions:	For Research Use only
Handling	
Storage:	-20 °C
Storage Comment:	Upon receipt, the positive control should be removed and stored at -20° or -80°C. The remainder
	of the kit can be stored for up to 6 months at 2-8°C from the date of shipment. Opened
	Microplate Wells or reagents may be stored for up to 1 month at 2° to 8°C. Return unused wells
	to the pouch containing desiccant pack, reseal along entire edge. Note: The kit can be used
	within one year if the whole kit is stored at -20°C upon receipt. Avoid repeated freeze-thaw
	cycles.
Expiry Date:	6 months