



[Go to Product page](#)

Datasheet for ABIN5526701
TR4 ELISA Kit

Overview

Quantity:	96 tests
Target:	TR4 (NR2C2)
Binding Specificity:	pSer412, total
Reactivity:	Human
Method Type:	Cell ELISA
Application:	ELISA

Product Details

Purpose:	Cell-Based ELISA Kit. This assay semi-quantitatively measures TAK1 phosphorylated at Serine-412 as well as total TAK1 in adherent cell lines.
Sample Type:	Adherent Cell Culture
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human TAK1 phosphorylated at site Serine-412 as well as total TAK1.
Characteristics:	<ul style="list-style-type: none">• Rapidly measure phosphorylated protein in adherent cell lines• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)• No sample lysis is needed• Compatible with a standard ELISA plate reader
Components:	<ul style="list-style-type: none">• Uncoated 96-well Strip Microplate• Wash Buffers

Product Details

- Fixing Solution
- Quenching Buffer
- Blocking Buffer
- Anti-phospho antibody
- Anti-pan antibody
- HRP-Conjugated Secondary Antibody
- TMB One-Step Substrate
- Stop Solution

Material not included:	<ul style="list-style-type: none">• Distilled or deionized water• 100 mL and 1 liter graduated cylinders• Tubes to prepare sample dilutions• Protease and Phosphatase inhibitors• Precision pipettes to deliver 2 μL to 1 mL volumes• Adjustable 1-25 mL pipettes for reagent preparation• Benchtop rocker or shaker• Microplate reader capable of measuring absorbance at 450 nm
------------------------	---

Target Details

Target:	TR4 (NR2C2)
Alternative Name:	TAK1 (NR2C2 Products)
Gene ID:	6885
UniProt:	O43318
Pathways:	TCR Signaling , Nuclear Receptor Transcription Pathway , Steroid Hormone Mediated Signaling Pathway , Regulation of Leukocyte Mediated Immunity , Positive Regulation of Immune Effector Process , Production of Molecular Mediator of Immune Response , Tube Formation , Toll-Like Receptors Cascades

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Plate:	Pre-coated
Protocol:	<ol style="list-style-type: none">1. Prepare all reagents and samples as instructed in the manual.2. Add 100 μL of sample or positive control to each well.3. Incubate 2.5 h at RT or O/N at 4 $^{\circ}$C.4. Add 100 μL of prepared primary antibody to each well.5. Incubate 1 h at RT.

Application Details

6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.
10. Add 50 µL of Stop Solution to each well.
11. Read at 450 nm immediately.

Restrictions: For Research Use only

Handling

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

Expiry Date: 6 months