

[Go to Product page](#)

## Datasheet for ABIN5526718

**WNK1 ELISA Kit**

## Overview

Quantity:	96 tests
Target:	WNK1
Binding Specificity:	pThr60, total
Reactivity:	Human
Method Type:	Cell ELISA
Application:	ELISA

## Product Details

Purpose:	Cell-Based ELISA Kit. This assay semi-quantitatively measures WNK1 phosphorylated at Threonine-60 as well as total WNK1 in adherent cell lines.
Sample Type:	Adherent Cell Culture
Analytical Method:	Semi-Quantitative
Detection Method:	Colorimetric
Specificity:	This ELISA kit recognizes Human WNK1 phosphorylated at site Threonine-60 as well as total WNK1.
Characteristics:	<ul style="list-style-type: none"><li>• Rapidly measure phosphorylated protein in adherent cell lines</li><li>• Simultaneously measure Phosphorylated protein and pan protein in one experiment (for normalization purpose)</li><li>• No sample lysis is needed</li><li>• Compatible with a standard ELISA plate reader</li></ul>
Components:	<ul style="list-style-type: none"><li>• Uncoated 96-well Strip Microplate</li></ul>

## Product Details

---

- Wash Buffers
- Fixing Solution
- Quenching Buffer
- Blocking Buffer
- Anti-phospho antibody
- Anti-pan antibody
- HRP-Conjugated Secondary Antibody
- TMB One-Step Substrate
- Stop Solution

Material not included:

- Distilled or deionized water
- 100 mL and 1 liter graduated cylinders
- Tubes to prepare sample dilutions
- Protease and Phosphatase inhibitors
- Precision pipettes to deliver 2 µL to 1 mL volumes
- Adjustable 1-25 mL pipettes for reagent preparation
- Benchtop rocker or shaker
- Microplate reader capable of measuring absorbance at 450 nm

## Target Details

---

Target: WNK1

Alternative Name: WNK1 ([WNK1 Products](#))

Gene ID: 65125

UniProt: [Q9H4A3](#)

## Application Details

---

Application Notes: Optimal working dilution should be determined by the investigator.

Plate: Pre-coated

Protocol:

1. Prepare all reagents and samples as instructed in the manual.
2. Add 100 µL of sample or positive control to each well.
3. Incubate 2.5 h at RT or O/N at 4 °C.
4. Add 100 µL of prepared primary antibody to each well.
5. Incubate 1 h at RT.
6. Add 100 µL of prepared 1X HRP-Streptavidin to each well.
7. Incubate 1 h at RT.
8. Add 100 µL of TMB One-Step Substrate Reagent to each well.
9. Incubate 30 min at RT.

## Application Details

---

- 10. Add 50 µL of Stop Solution to each well.
- 11. Read at 450 nm immediately.

---

Restrictions: For Research Use only

## Handling

---

Storage: -20 °C

Storage Comment: Upon receipt, the kit should be stored at -20 °C. Please use within 6 months from the date of shipment. After initial use, Wash Buffer Concentrate (Item B), Assay Diluent (Item E), TMB One-Step Substrate Reagent (Item H), HRP-Streptavidin (Item G), Stop Solution (Item I) and Cell Lysate Buffer (Item J) should be stored at 4 °C to avoid repeated freeze-thaw cycles. Return unused wells to the pouch containing desiccant pack, reseal along entire edge and store at -20 °C. Reconstituted Positive Control (Item K) should be stored at -70 °C.

---

Expiry Date: 6 months